

Dental Digest

December 1953


IN THIS ISSUE

- Chemical and Mechanical Aids in
Intraoral Roentgenography 530
- Chemistry In Health and
Disease—Part Four 536
- Medical Conditions Complicating
Oral Surgery 544
- Penicillin Used in
Odontologic Cases 545
- Clinical and Laboratory
Investigations 548
- Pharmacology and the Biologic
Sciences 550
- Annual Index 553
- Pathologic and Neurologic
Inclusion 562
- Cardiac Patient as a
Surgical Risk 565

(A complete table of contents
appears on page 529)

Cover illustration—Shapiro article
Page 530

UNIVERSITY
OF MICHIGAN
DEC 24 1953
DENTISTRY
LIBRARY



OUTSTANDING...

IN BOTH MATERIALS

VERI-CHROME PORCELAIN AND VERIDENT PLASTIC

In Veri-chrome Natural Tooth Colors
USE OR SPECIFY FIVE-PHASE ANTERIORS IN PORCELAIN OR
VERIDENT PLASTIC WITH THE VERI-CHROME COLOR GUIDE




The most lifelike porcelain and plastic teeth ever made...

VERI-CHROME PORCELAIN: The famous time-tested Veri-chrome Porcelain provides the depth, refraction and translucency of natural tooth structure in a porcelain that is outstanding for its strength as well as its "living" character.

VERIDENT PLASTIC: The only plastic teeth that "come clean" from investment and retain their original lustrous surfaces and color without marring or discoloring tooth surfaces. The reason for this is that Verident Plastic Teeth have an extra safety-margin of resistance to heat, solvents, crazing, brush-flaming and surface attack, *built-in the material.*

- VERIDENT PLASTIC TEETH ARE IDENTIFIED BY A RED DOT ON THE LAP OF EACH TOOTH.
- FIVE-PHASE ANTERIORS IN VERI-CHROME PORCELAIN ARE IDENTIFIED BY A PENTAGON ON THE LAP OF EACH TOOTH.



FREE This helpful literature is available upon request:

1. Illustrated Mold Chart
2. 60 Personalized Five-Phase Anterior Denture Arrangements
3. 32-page Book: Varied Labial Surfaces and Their Relation To Scientific Tooth Selection

FIVE-PHASE

A N T E R I O R S

I. Sh
New
1916
with
and
Veter
Alban
is Fr
ORAL

MELV
Mich
foun
throp
publi
artic
lism
ratio
Colle
sistan
prese
of an
AND

HARR
sity,
use o
fistry
two
biotic

DECE

Dental Digest

Registered in U.S. Patent Office

DECEMBER 1953

Film and Mechanical Aids in Intraoral Roentgenography <i>I. Shapiro, D.D.S.</i>	530
Civil Malpractice (An Abstract)	535
The Chemistry of Disease (An Abstract)	535
Body Chemistry In Health and Disease—Part Four <i>Melvin E. Page, D.D.S., and D. L. Brooks, A.B.</i>	536
Medical Conditions Complicating Oral Surgery (An Abstract) <i>Benjamin Eliasoph, M.D.</i>	544
Terramycin Used in Endodontic Cases <i>Harry Maeth, D.D.S.</i>	545
The Editor's Page	552
Contra-Angles	556
Clinical and Laboratory Suggestions	548
1. A Plaster Splint for Putrescent Pulp. 2. Increasing the Rotating Speed of a Bur. 3. A Suregrip Sprue. 4. Patching an Alginate Impression. 5. A Simple Mount for X-rays. 6. A Hanger for X-ray Films.	
Medicine and the Biologic Sciences	550
This Is a Human Body (An Abstract)	551
Annual Index	553
Physiologic and Neurologic Occlusion (An Abstract)	562
The Cardiac Patient as a Surgical Risk (An Abstract)	565

About Our**CONTRIBUTORS**

I. SHAPIRO, Colonel USAFR (DC) (D.D.S., New York University College of Dentistry, 1916) has been associated professionally with the Veterans Administration since 1946 and is now Chief of Dental Service of the Veterans Administration Regional Office in Albany, New York. Colonel Shapiro's article is **FILM AND MECHANICAL AIDS IN INTRA-ORAL ROENTGENOGRAPHY**.

MELVIN E. PAGE, D.D.S. (UNIVERSITY OF MICHIGAN, COLLEGE OF DENTISTRY, 1919), founder of the American Society of Anthropometric Medicine and Nutrition, has published two books and a number of articles on the various aspects of metabolism and dental disease. With the collaboration of **D. L. BROOKS, A.B.** (Wheaton College, Norton, Massachusetts, 1931), assistant in dental medicine, Doctor Page presents the fourth installment of a series of articles, **BODY CHEMISTRY IN HEALTH AND DISEASE**.

HARRY MAETH, D.D.S. (Columbia University, 1925) follows his initial report on the use of the antibiotic, terramycin, in dentistry with an illustrated progress report of two endodontic cases in which the antibiotic was successfully used.

EDWARD J. RYAN, B.S., D.D.S., Editor**WANDA T. PICKARD, B.A., Assistant Editor**

708 Church Street, Evanston, Illinois

Copyright, 1953, by Dental Digest, Inc. See page 528 for subscription data, etc.
The magazine is mailed on the fifteenth of the month of issue.

Film and Mechanical Aids

in INTRAORAL ROENTGENOGRAPHY*

I. SHAPIRO, D.D.S., Albany, New York

DIGEST

This discussion of some of the problems encountered in intraoral dental roentgenography includes a detailed description of an appliance and film devised to eliminate distortion in dental x-rays. Step-by-step directions for procedure in using the device are given.

Distortion in Dental X-Rays

Bending of the dental x-ray film during exposure with consequent image distortion should be avoided as the basis of any technique the aim of which is the closest approach possible

to an exact anatomic reproduction of the teeth or area in question. Several conditions contribute to distortion.

Digital Pressure—Curvature of the arch and of the roof of the mouth makes elimination of distortion impossible as long as the patient must retain in position by digital pressure the oversized flexible films available for use at present.

Folded Film a Cause of Distortion—To overcome the inevitable poor results, particularly in the more restricted anterior part of the mouth, many operators resort to folding the film, which reduces the size to facilitate placing in position, but increases rigidity of the film.

One Accurate Representation Secured—Although the average exposure will contain three and sometimes four teeth, actually, because of dis-


tortion, only one (that in the center as a rule) can be relied upon as an adequate representation of the tooth in question. In the others, distortion increases proportionately as the periphery of the film is reached.

Exact Film Position Difficult to Obtain—A narrow rigid film reduced to the minimum in overall length for a single tooth exposure,¹ represents the nearest approach to the ideal. Obtaining the proper position by visual means, however, presents a problem, since the vertical center line of the film must coincide with the buccolingual longitudinal plane through the long axis of the tooth in order to secure its full outline within the confines of the film. A mechanical aid to accomplish proper placement would be of value.

Favored Technique—Most opera-

*Reviewed in the Veterans Administration and published with the approval of the Chief Medical Director. The statements and conclusions in this article are the results of the author's personal investigation and do not necessarily reflect the opinion or policy of the Veterans Administration.

¹Pollio, Joseph A.: *Fundamental Principles of Alveolo-Dental Radiology*, Brooklyn, New York, Dental Items of Interest Publishing Co., 1933, pp. 89-107.



1. The two main divisions of the device are shown. The section containing the bite-block and slotted shaft for the film packet (which lies in the mouth) has three types of film displayed behind it: (1) a single tooth at the top used mainly for anteriors, (2) a wider one to include two teeth for bicusps and molars, and (3) the bite film at the bot-

tom which takes in only one maxilla. The latter, if anything, tends toward greater accuracy than the type now in use.

2. A quarter turn after insertion locks the mouth section (buccal aspect) into the adjoining or indicator section, making the implement ready for use.

tors find it less difficult to secure the proper vertical angle by using the bisecting angle technique. The horizontal angle, however, requires greater accuracy as the crowns of the teeth normally are in contact proximally and the slightest deviation results in an overlapping of these surfaces, obliterating the area.² Procuring both angles readily at the same time would be ideal. This would necessitate accurate placing of the x-ray tube.

Means of Securing Proper Vertical and Horizontal Angles

To accomplish satisfactory results the following three conditions must be present:


1. A rigid film packet reduced to the minimum in size to permit of latitude in placement in confining areas, and definite in outline to ensure its being centered in the device.
2. A mechanical device which would readily secure the packet in accurate position so that guidance of the central ray through the center of the film would be automatic while conforming to the basic principles of the bisecting angle or right-angle technique.

3. A means of ensuring the proper alignment of the x-ray tube to the film so that the correct vertical as well as horizontal angles are secured simultaneously.

Achievement of the requirements listed must be simple and uncomplicated for satisfactory application of the entire procedure:

Introduction of Indicator Bar

A device has been developed which will accomplish the desired results using the bisecting angle technique. While the device can be modified for the right angle (long cone) technique,⁴ in its present state addition of a bite-block extension makes the appliance readily available for use when x-raying the posterior teeth;



3. The top view shows the relationship of the film packet "F" to bite-block "B". The center line of the film is continuous with that (guide line "G") of the bite-block. The distance between "G" and indicator bar "IB" equals that between central ray "R" and extension bar "EB". The target film distance "TFD" is always constant.

that is, bicuspid and molars. A combination of both techniques, therefore, is feasible in full mouth exposures. Another attachment which fits over the indicator bar converts it for use in stereoroentgenography. No attempt at refinement of the device has been considered at this time, the object being solely to demonstrate its usefulness and practical aspects.

Description of Device

The rigid plastic framework of the film packet which prevents bending has a definite extension at the bottom with converging sides, each equidistant from the middle S (Fig. 2). This design ensures centering the film when the packet is inserted in the device. Regardless of the length or width, a line drawn through the center of the film will coincide with the horizontal guide line which divides the bite-block into halves (Fig. 3). The appliance is visualized and described in Figures 1 to 13.

Additional Provisions

In the case of upper first bicuspid, provisions could be made to permit

rotating the indicator bar about 25 degrees, in a horizontal plane, around the upright "U" as an axis, thus shifting the x-ray cone so that exposure will reveal two roots where these are present rather than one normally resulting from superimposition of the buccal on the lingual root. Maintaining the patient's head so that the plane of occlusion is parallel to the floor while the sagittal plane is perpendicular at the same time, would not be required.

Possible to Eliminate Preliminary Steps—The conventional head position must be used, however, because of the universal type of x-ray tube suspension which limits its movements to a vertical and horizontal plane. An improvement would be a change to a ball and socket arrangement permitting adjustment in virtually any position, or the inclusion of a vertical swivel joint, at the point of suspension above the head, would eliminate this preliminary step entirely.






²Fitzgerald, Gordon M.: Supplementary Roentgenographic Examinations, N.Y. State Dent. J. 15:449-456 (Oct.) 1949.

⁴McCormack, Donald W.: Intraoral Roentgenology, DENTAL DIGEST 57:106-114 (Mar.) 1951.

¹Ennis, LeRoy M.: Dental Roentgenology, Philadelphia, Lea & Febiger, 1942, pp. 75-117.

²McCall, J. O., and Wald, S. S.: Clinical Dental Roentgenology, Technique and Interpretation, ed. 2, Philadelphia, W. B. Saunders Company, 1940, pp. 88, 89.

³Fitzgerald, Gordon M.: Dental Roentgenography II: Vertical Angulation Film Placement and the Extended Object, Film Distance, JADA 34:160-170 (Feb. 1) 1947.



4. The film is inserted into the mouth. Moving indicator "I" in the same direction, by pressure of thumbnail on the left nut, revolves the shaft so that the film assumes a horizontal position which facilitates its entry into the mouth. Black line "F" represents the edge of the film packet.





5. The guide line between the bicuspid and molar automatically places the center of the film in the same position. After closing the teeth, release of the indicator permits the shaft to rotate (by means of an internal spring) toward the vertical, with the film packet coming to rest as it comes in contact with the tooth and lingual tissues as shown in Figures 6 and 7.

6. The upright segment "U" (extending from indicator "I") by means of a mechanical arrangement, automatically bisects the angle formed by the planes of the film and the long axis of the tooth. The indicator bar "IB" at a right angle to the upright segment "U", denotes the proper direction to be taken by the central ray.

7. Alining the extension bar on the yoke which fits over the x-ray cone with the indicator bar will simultaneously result in the proper vertical and horizontal angles. The fixed length of the indicator and extension bars ensures the same target distance at all times.

8. The bite-block extension "BE" for right angle or long tube technique in the posterior part of the mouth permits the position of the film in use as shown in Figure 9.

9. The film in position parallel to the long axis of the teeth in or beyond the median line with upright "U" parallel to it. Alinement of the indicator and extension bars will result in the central ray passing at a right angle through the center of the film. This situation is based on the premise that the occlusal surface of a tooth (for this purpose) is at a right angle to its long axis. Since film and tooth are at right angles to the surface of the bite-block, it follows that they are parallel to each other as well as to the upright "U."




10. Stereoroentgenography attachment "SE" which fits over the indicator bar.

11. First exposure with the bars aligned in the left position. The movable attachment "M" is shown.

12. The second film and exposure in the correct position are demonstrated. The movable end of the attachment "M" is swung out of the way. For additional accuracy, rather

than relying entirely on the bite-block guide line, baseplate wax or compound luted to the under surface of the bite-block and used as an index of the teeth will ensure exactly the same position when making both exposures.

13. A composite view showing how attachment design (calibrated) produces an intertarget (interpupillary) distance of 65 millimeters.



14. Shows a full mouth series where an individual exposure is made for each of the anterior teeth.

over
ously
fixed
same

long
rmits

teeth
el to
result
enter
at the
right
right
they
"U."

X-Rays of Individual Teeth Possible—Although the most desirable results would be obtained by x-raying each tooth individually, this method is impractical⁶ because of the difficulties involved. Use of the procedure advocated, however, makes individual x-rays possible; this method was used in obtaining the radiographs shown, from the upper left bicuspid to the right cuspid inclusive, as well as the lower anterior teeth. The method could be used for all the teeth (Fig. 14).

Attachment for Use in Stereoroentgenography

Because of the difficulties involved in attempting stereoroentgenography with the present dental x-ray equipment, this valuable adjunct has become almost a lost art.⁵ The use of the attachment illustrated in Figures 10, 11, and 12, it is believed, would encourage more general use.³


Bite-block Cover Provided—A thin individual plastic bite-block cover is planned. This device, containing the tooth registration in compound (shown in Figure 12), is an aid in placing both films accurately in the same position. The bite-block cover can also be used and retained for future use in all cases where a follow-up check is contemplated.

Method Advocated

Standard type periapical films were cut to sizes $1\frac{1}{4}$ inches by $1\frac{1}{16}$ inches for bicuspid and molars and $1\frac{3}{8}$ inches by $\frac{5}{8}$ inch for cuspids. Although strips $\frac{5}{8}$ inch wide (for incisors) were obtained by cutting across the narrower film dimension to facilitate their use, those for the cuspids could be used as well, thus confining the assortment of film in the mouth to two.

Full Size Packet Used—In the lower third molar region the full size packet was used, since an adjustment not incorporated at the present time precluded the use of the method described for bicuspid and molars in extending to the retromolar area to complete the full mouth series.

Bending of Film Prevented—A thin sheet of rigid steel plate, similar in



15. An upper bicuspid view with the exposures made and the films mounted for stereoscopic study.

16. An impacted upper cuspid with the exposure made and the films mounted for stereoscopic study.

outline to the film designs illustrated in Figures 1 and 2, was attached by means of scotch tape to the cut packet lingually in each instance before placing it in the device. This step precluded bending of the film thereby simulating the effect obtained by use of the rigid plastic framework recommended for the ready-made packet.

Increase in Length—The initial results show a slight increase in film length indicated for the cuspid teeth. This can be provided for, considering the fact that the cut length was $1\frac{1}{4}$ inches in contrast to the long dimensions of $1\frac{5}{8}$ inches of the films now in use.

Stereoscopic Views for Additional Study—Both upper first bicuspid show a double root formation. For further study, stereoscopic views of the right one were taken. The results may be seen by examining the mount through a hand stereoscope (Fig. 15).

An additional view of an impacted cuspid is shown to reveal the exact location of the unerupted tooth (Fig. 16).

Stereoscopic Pictures Feasible—Although the Standard Dental X-Ray Unit without a stereoshift is not constructed to permit the exact movements necessary in stereoroentgenography and the films were cut by hand, while the device in its initial stage, as well as the procedure, was somewhat crude, the results obtained nevertheless demonstrate the practical aspects of obtaining stereoscopic views readily by the procedure described.

Additional Effort Warranted by Results—Obtaining occlusal roentgenograms involves less than the periapical because the film holder (not shown), acting as a receptacle, is not removed from the mouth during the changing of the films. The additional

information gained from these views, compared to the ordinary single flat plane, warrants the small additional expenditure of time and effort required.

Central Ray Controlled—The ability virtually to pinpoint focusing of the central ray permits restricting the circle of useful radiation to the smallest practical diameter with the ultimate effect of confining it to the approximate size of the film. Unneces-

sary exposure of the patient is avoided, and as stressed, potential danger to the operator by direct and stray radiation is avoided.

Summary

1. A mechanical device is presented which will indicate the proper positioning of the dental x-ray cone during exposure, using the bisecting angle technique with an attachment to include the right angle technique for bicusps and molars.

2. An additional means is provided to facilitate alinement of the x-ray tube to secure proper vertical and horizontal angles simultaneously.

3. The procedure involves the use of a rigid periapical and bitewing dental film packet, definite in outline design, and reduced to the minimum in size to facilitate placement.

4. The appliance is also adaptable for stereoroentgenography.

560-A New Scotland Avenue.

We Can't Pay You, But—

NO DENTAL AUTHOR can ever be paid for a valuable technical or scientific article. The value of such material is above a monetary basis. In the preparation of a technical article, however, an author often spends money for drawings, photographs, models,

or graphs. We should like to help defray some of these expenses.

Until further notice, DENTAL DIGEST will allow \$25.00 toward the cost of the illustrations provided by the author of every article accepted.

If you have a constructive idea, an

innovation, a new result of tried and proved experiment, put it down in writing, illustrate it, and send the material to: DENTAL DIGEST, 708 Church Street, Evanston, Illinois.

We can make suitable black-and-white cuts from Kodachrome transparencies.

We hope that you will accept this invitation!

Civil Malpractice

Important Points in Preventing Malpractice Claims

1. Handle cases only within your professional qualifications; call in consultants whenever necessary.

2. Do not be overoptimistic in prognosis.

3. Always obtain adequate consent.

4. Always keep adequate re-

cords for your own as well as your patient's protection.

5. Always have x-rays taken of bone and joint injuries. Use laboratory test, even though simple, when indicated.

6. Make frequent enough calls and give sufficiently precise instructions so that there can be no misunderstanding. Be sure you write your prescriptions legibly.

7. Be sure that your patients

are covered by medical attention in the event of your absence.

8. Know what to do when the patient does not follow instructions or discontinues treatment.

9. Do not gossip about your cases.

10. Do not criticize your professional colleagues.

From *Medical Clinics of North America* 37:No. 5, 1575 (September) 1953.

The Chemistry of Disease

MORE SHOULD be known of the chemistry of disease. When it is, physicians will not be content to accept the traditional view that when a person is not well, or as one ages, he must expect to lose strength, keenness of vision, and alertness and to look and act "old." The physician can never get

away from the challenge to think. There will be no automatic or mechanical diagnosis in the near future. The physician who has an uncanny sensitiveness associated with a capacity for memory and experience is needed more in this field than in any other. He must have in mind that there is nothing

constant in the human body. The nutrients ebb and flow into the cells. They all bear potentials for change. In this ever-shifting, ever-renewing of the constituents of the body, the cells must be renewed or there will be deterioration. These forces shape the patient's destiny.

—Tom D. Spies, M.D.: *JAMA* 153:185-193 (Sept. 19) 1953.

BODY CHEMISTRY

in Health and Disease—Part Four

MELVIN E. PAGE, D.D.S., and D. L. BROOKS, A.B., St. Petersburg, Florida

DIGEST

The endocrine content of the blood, although minute, is not insignificant; it is capable of maintaining life. The assumption is therefore justified that minute quantities of endocrine substances can also be used to obtain health. Systemic factors related to oral health should receive the attention of dentists: the patient presents more than an oral problem and if he is to receive maximum health service he should be considered as a total organismal entity.

Graph interpretation and the response of calcium-phosphorus blood levels to minute amounts of endocrines demonstrate hypo and hyperfunctioning glandular activity for the thyroid, posterior pituitary, and anterior pituitary glands. Success in changing calcium-phosphorus blood levels by these means is dependent upon the right endocrine in the right amount for the particular person.

The outstanding physical characteristics of each gland in hyper-

or hypo-function are noted with comment upon changes in blood sugar levels, specific gravity of the urine, and blood pressure. The conclusion is that the thyroid gland is opposed by the insulin-producing cells of the pancreas, the posterior pituitary gland is opposed by the adrenal cortex, and the anterior pituitary gland is opposed by the pancreas and/or the sex hormone glands.

Clues to endocrine dosage are (1) the degree of graph divergence from normal, (2) age, and (3) correlation of various blood indexes. The intent of this therapeutic approach is to supplement glandular production only to the extent of the gland's own limitation. To supplement in excess of need or normal production would defeat the purpose of establishing an efficient body chemistry. Too little is known about the body's daily production of endocrines, but indexes of efficient body chemistry where norms have already been established can be accepted as indicators of progress.

Purpose of Treatment and Dosage

The significant difference between endocrine treatment based on the approach described in these articles and endocrinology as practiced by physicians is: In medicine endocrines are used as a rule in large doses to achieve a pharmacologic response; in this approach endocrines are applied

in minute doses to make up deficiencies in the subject's glandular output. The medical approach produces an immediate response to overcome an acute situation; this approach produces a gradual increase in chemical efficiency to overcome an inherited or chronic inefficient body chemistry sometimes called a state of low resistance. The purpose of treatment differs

to the same extent that dosage differs.

Ingle and Baker¹ distinguish between the *physiologic* dosage (the amount does not exceed the normal secretory output of a gland) and the *pharmacologic* dosage (the amount is greater than that normally secreted by a gland). In their words: "The biologic response to high dosage of any drug or hormone tends to decrease with time due to the general phenomenon of adaptation . . . There is a growing tendency to employ the smallest doses of hormone which will initiate a remission, not always complete, and to subsequently reduce the dose to achieve the best possible suppression of symptoms consistent with avoidance of significant complications."

Endocrine Dosage Infinitesimal—Generally speaking, the medical profession would consider the endocrine dosage advocated in this technique so small as to be insignificant. Even biochemists refer to the endocrine content of the blood as so small as to be insignificant. The endocrine elements in the blood are in fact so minute that under present methods of chemical analysis they are almost incapable of fractionization.

Life Maintained by Blood Content—It is, nevertheless, an undisputed fact that the blood is the transporting medium of endocrine secretions within the body. Without endocrine substances death occurs.

Health and Ill Health—If such minute amounts of endocrines make the difference between life and death it is reasonable to conceive of minute amounts making the difference between health and illness.


¹Ingle, Dwight J., and Baker, Burton L.: *Physiological and Therapeutic Effects of Corticotropin (ACTH) and Cortisone*, Springfield, Illinois, Charles C Thomas, Publisher, 1953, p. 81.

BIOCHEMICAL INDEX

BLACK LINE REPRESENTS PHOSPHORUS LEVEL — RED LINE CALCIUM LEVEL

DATE	B.S.	B.P.	U.S.G.	CA.	P:	1	2	3	4	5	6	7	8	9	10	11	12															
1st Blood						WC	R	C	Hgb	C.V.	ph	Dif	Eos	T.N	Seg	Stab	Juv	Myel	Lym	Mono	Heag	C	II	V.I	Ani	Poi	Orth	Achr	Ret	Plate	Sed	
2/12/53	111					10.2					7.5																				16	
2nd - In absence of sugar																																
2/19/53	100					10					3.5																					46
						Medication					8.9	7.5																				

Length right forearm	Quotients	Length right leg
9.9	1.688	16.7
Wrist	6.15 A 1.610	9.9 Ankle
one-fourth	6.15 B 1.414	8.7 one-fourth
one-half	7.5 C 1.565	11.75 one-half
three-fourth	8.75 D 1.366	11.95 three-fourth
Elbow	9.6 E 1.544	14.8 Knee



Copyright 1950
By MELVIN E. PAGE, D.D.S.

Factors to be Discussed

Changes in Blood Indexes—In the following discussion the effect upon calcium-phosphorus levels in the blood will be of particular interest because of its importance in a systemic understanding of dental ills. Regardless of the focus and limitation of his activity the dentist should have an understanding and appreciation of underlying systemic factors which affect the patient as a whole being.

The Patient a Total Organism—The patient who sits in a dental chair does not present merely an oral problem. He is first of all a human being composed of many indivisible parts, emotional reactions, and mental attitudes. He must be understood, approached, and treated as a complete entity, a total organism, if he is to receive the ultimate in health service. This is the unitary concept of therapy.

Method of Presentation—Before demonstrating a complete case history, studies will be made involving one gland alone. The graph indications for endocrine choice will be discussed. Blood tests prior to treatment will be recorded for comparison with

blood analyses obtained following treatment. Studies of hypofunctioning glands will be followed by study of the same gland in hyperfunction.

Corrected Diet-Definition—It should be understood that unless otherwise specified the patients represented in this article were following a corrected diet at the time blood analyses were taken. *Corrected diet means no sugar in any form, no fruit juice, no white flour, no milk, no alcohol, and immediately prior to a blood test, not even any whole fruit, and no coffee on the day of the test.*

Analysis of Figure 31

The outstanding physical characteristic of a person with a hypofunctioning thyroid is heaviness of the leg just above the ankle, the first 1/4 circumferential leg measurement.

Hypofunctioning Thyroid Shown—The patient in this case (Fig. 31) is a woman with a hypofunctioning thyroid. The starting point of the graph, it will be noted, is 1.610. This is higher than normal; therefore, the reading will be simplified if a normal graph is superimposed upon the chart. The starting points, for both the specific

graph and the normal, coincide. It will be recognized that the dotted line coming above the normal solid line indicates a hypofunctioning thyroid which was described in Installment Two of this series of articles.

Results of Blood Analysis—The blood analysis (Fig. 31), taken prior to the use of thyroid extract, shows a blood sugar of 111, calcium 10.2, phosphorus 3.07, hemoglobin 85, cell volume 43.6, blood pH 7.5, sedimentation rate 16, according to the Clay-Adams method of determination. (Normal sedimentation rates: men, 0-8; women, 0-10, 12-14 during menses.) Since phosphorus utilizes calcium at the rate of 2.5 times itself, multiply the phosphorus of 3.07 by 2.5. The answer is 7.67. This is the usable compound level of calcium and phosphorus.

Excess Calcium Present—In the biochemical index note that the solid calcium line extends beyond the pencil point (7.67) to 10.2 (the calcium level at the time of this analysis). This means that there is an excess of 2.5 milligrams of free, uncombined calcium. This excess is consistent with the hypofunctioning thyroid graph

and the low compound level is consistent with the occurrence of yearly cavities.

*Additional Factors Considered—*The blood sugar level of 111 is indicative of carbohydrate intolerance. The blood pH is alkaline, being 7.5 instead of the normal 7.4. Sedimentation rate should be 0-10 (12-14 during menses) for a woman but in this instance is 16. Hemoglobin is slightly low, being 85. Cell volume 43.6 is virtually perfect, 43 being normal for a woman.

Endocrine Therapy Applied—One-fiftieth (1/50) grain of thyroid extract was given daily for seven days. The blood analysis was then repeated with the results charted in Figure 31 (2/19/53): Blood sugar 100, calcium 10, phosphorus 3.5, hemoglobin 92, cell volume 38.7, blood pH 7.5, sedimentation rate 16.

Normal Blood Sugar—The blood sugar in this subject is now a perfect 100. In this approach this is normal and anything above or below is abnormal although medical textbooks quote normal blood sugar levels at anywhere from 80 to 120. Probably this is true of the *average* person on the usual American diet: nevertheless,

in a person eating a corrected diet, and with an efficient body chemistry the mechanism controlling carbohydrate metabolism should function with such perfection that the blood sugar remains at 100.

Improvement Noted—The following ratios exhibit definite improvement following thyroid extract therapy:

Calcium-Phosphorus Rate Improved: Not only has the blood sugar improved but also the calcium-phosphorus ratio. Under the biochemical index, if a dot is placed at the usable compound level, 8.75 (multiply the phosphorus 3.5 by 2.5) an increase in the *usable* compound level of calcium-phosphorus will be noted. This is a step in the right direction for improvement of dental health.

Reduction in Excess Calcium: At the same time that the usable compound level increases, there is a reduction in the amount of excess calcium. At first there were 2.5 milligrams of excess calcium; now there is less than 1.5 milligrams of excess calcium. Improvement of blood sugar, calcium-phosphorus ratio, and compound levels was due to the use of thyroid extract. The choice of thyroid extract

was dictated by the endocrine interpretation of the graphed body measurements.

*Basis of Successful Treatment—*Success in changing calcium-phosphorus levels by endocrine means is dependent upon choosing the right endocrines in the right amount, for the right person. This rule should be kept in mind while studying these reports.

In the words of Walter Cannon:² "Treatment should serve in a natural manner to perform a natural function that has broken down. Use a physiological factor for a physiological defect."

Analysis of Figure 32

The study presented in Figure 32 is a case of hyperthyroidism in a male in contrast to the previous hypothyroid female.

Characteristics of Hyperthyroidism—The outstanding physical characteristic of a hyperthyroid person is the slimness of the leg just above the ankle bone (Fig. 32). The starting point on the graph for this male patient is 1.442. When the patient's

²Cannon, Walter B.: *The Wisdom of the Body*, New York, W. W. Norton & Company, Inc., 1932, p. 242.

BIOCHEMICAL INDEX

BLACK LINE REPRESENTS PHOSPHORUS LEVEL — RED LINE CALCIUM LEVEL

[illegible]

graph is plotted, the dotted line representing his measurements falls too low on the vertical line BB'. This is indicative of a hyperfunctioning thyroid.

Results of Blood Analysis—A blood analysis prior to treatment (1/12/52) but on corrected diet, shows a blood sugar of 105, calcium 8.8, phosphorus 2.9, hemoglobin 93, cell volume 57.1, blood pH 7.4, sedimentation rate 6, blood pressure 176/110.

Factors Concerned in Blood Pressure—It is commonly known that the average blood pressure increases with age. It is not so well known that a normal blood pressure remains at the same level throughout adult life. The normal blood pressure is 113-120 over 70. The upper pressure is the systolic; the lower pressure is the diastolic. Most people know about their systolic blood pressure but few appreciate that the diastolic is the more important in cases of hypertension. In this instance, the 110 diastolic is excessive.

Effects of Related Glandular Deficiencies—A high phosphorus blood level might have been expected in the hyperthyroid person in this case. This does not necessarily occur, however, when other glandular inadequacies are also involved. To estimate the degree of imbalance in the blood picture, multiply the phosphorus of 2.9 by 2.5. The answer of 7.25 provides the usable compound level. The difference between this 7.25 and the calcium blood level of 8.8 is 1.55 milligrams, the amount of excess calcium.

Overactivity a Treatment Problem—In the previous case (Fig. 31) thyroid extract was used because the patient's own gland was underfunctioning. In this case (Fig. 32) there is a hyperfunctioning or an overactive gland. This gland is either making too much thyroxine or an abnormal substance. The problem, therefore, is not to make up for a deficiency in output, but to inhibit overactivity.

Endocrine Therapy Applied—It has been found through experience that thyroid substance is opposed, at least in part, by insulin. Consequently, the patient was given 4 units of 40-unit protamine zinc insulin once daily for six days. This dosage which would be

inadequate for the treatment of diabetes is nevertheless adequate when used for alteration of calcium-phosphorus blood levels.

Results of Initial Therapy—After taking 4 units of insulin once daily for six days, another blood analysis produced the following results:

Blood sugar 98, calcium 9.6, phosphorus 3.5, hemoglobin 100, cell volume 50, blood pH 7.4, sedimentation rate 10. Multiplying the phosphorus of 3.5 by 2.5 again, a usable compound level of 8.75 is demonstrated.

Excess Calcium Reduced—There is now less than 1 milligram of excess calcium in contrast to the original 1.5 milligrams of unused calcium.

Additional Favorable Alterations—Note also that the hemoglobin has risen to 100. The cell volume has dropped from 57.1 to 50 (45 is normal for a man). The pH of the blood remained at a normal of 7.4. The sedimentation rate has risen slightly, although it is still within normal limits. The blood sugar has changed from a little above normal to slightly below normal. The blood pressure, both diastolic and systolic, has dropped.

Cumulative Effect of Changes Significant—All these changes are small but cumulatively they are significant. All are indexes of increasing efficiency of body chemistry, increasing resistance to oral degeneration, and other forms of degeneration.

Comment

The effect has been demonstrated of two different endocrine substances upon calcium-phosphorus blood levels (thyroid and insulin). Opposing endocrine substances have been used for opposing endocrine patterns with improvement of calcium-phosphorus ratios in both instances.

Analysis of Figure 33

In this graph (Fig. 33) the high midpoint (the C-C' line) is indicative of a hypofunctioning posterior pituitary gland. The middle peak is higher than the normal. The outstanding physical characteristic for a hypofunctioning posterior pituitary is found about midway between the an-

kle and the knee. There is a marked curve at the calf of the leg at this point.

Note—It must be kept in mind that in this series of studies a deficiency or hyperactivity of one gland is being demonstrated at a time. Other glandular dyscrasias may exist in the same case, and often do, but for explanatory purposes attention is directed to only one.

Blood Analysis—On a corrected diet the blood analysis in this case shows a normal blood sugar of 100, a normal hemoglobin of 100, and a normal female blood cell volume of 44.4.

Divergencies from Normal—The ratio of calcium, 10.4, to the phosphorus is 3.25. The usable compound level is 8.00, leaving an excess of 2.5 milligrams of calcium. The blood pH is slightly high (7.45) and the sedimentation rate is at the abnormal level of 26. Blood pressure is 150/90, and the average specific gravity of the urine runs below normal.


Hypertension in Hypoposterior Pituitary Conditions—It is not unusual to find hypertension in people with hypoposterior pituitary glands. It is usual to find a lessening of the hypertension with correction of the deficiency in posterior pituitary function.

Kidney Mechanism—The posterior pituitary gland seems not only to have an indirect influence on blood pressure, but also upon the specific gravity of the urine. The kidneys take from the blood approximately a barrel of water daily. A barrel of water would be an inconvenience both in intake and output. In this situation nature is a successful conservationist. All fluid intake is circulated through Bowman's capsules of the kidneys. In passing through the kidney tubules much liquid is reclaimed for reuse. A relatively small amount of liquid is utilized to carry out unwanted waste material in the urinary flow. The amount of daily glomerular filtrate is thought to be 100 times the amount of urine excreted.

Method of Estimating Kidney Efficiency—The efficiency of the kidney mechanism is registered in the density of the outgoing urine: (1) A low spe-

BIOCHEMICAL INDEX

BLACK LINE REPRESENTS PHOSPHORUS LEVEL — RED LINE CALCIUM LEVEL



•33•

Copyright 1950
By MELVIN E. PAGE, D.D.S.

cific gravity of urine means low density and inadequate water conservation, and (2) the normal specific gravity reading should be 1.022 both morning and night.

Specific Gravity Level Improved—(1) If the figures indicating specific gravity obtained both night and morning for several days have a low average, and (2) if the graphed measurements indicate hypofunction of the posterior pituitary, the specific gravity levels will tend to improve with the use of the correct amount of posterior pituitary substance for that person.

Therapy Administered—The patient in this case (Fig. 33) was given 1/400 grain of posterior pituitary extract daily for five days. A blood specimen taken at the end of that interval (11/3/52) showed an undesirable sugar level of 125 which indicates that posterior pituitary supplement is not the only requirement in this case.

Improvement Demonstrated—Calcium-phosphorus ratios improved markedly with therapy. The calcium blood level was 10.2 and the phosphorus blood level 3.8, producing a compound level of 9.5. This left less than 1 milligram of excess calcium, as op-

posed to the original 2.5 milligrams of excess calcium.

Analysis of Figure 34

The graph of the patient in this case (Fig. 34), that of a hyperfunctioning posterior pituitary in a female, shows the midpoint flattened and falling below the normal.

Typical Physical Characteristic—Straightness of the leg between the ankle and knee is the chief physical characteristic of the person in this graph. The midcalf has little or no bulge or curve.

Results of Initial Blood Analysis—The first blood test on the corrected diet shows a blood sugar of 100, calcium 10.4, phosphorus 3.7, cell volume 44.4, blood pH 7.59, sedimentation rate 10. The phosphorus of 3.7 multiplied by 2.5 gives a compound level of 9.25. There is one milligram of excess calcium.

Minute Amount of Hormone Administered—The patient was given 1/10 grain equivalent of adrenal cortex daily. The effect of this dosage is shown (11/29/42) in a blood sugar reading of 114, calcium 9, phosphorus 3.6, cell volume 38.7, blood pH 7.4, sedimentation rate 10.

Increase in Blood Sugar Level Interpreted—There is no excess of calcium. A perfect balance of calcium-phosphorus is present, but the increase in the blood sugar level would indicate (1) that the dosage of adrenal cortex is too strong, or (2) that other glandular supplementation must be used concurrently with the adrenal cortex.

Method to Inhibit Overactive Posterior Pituitary—Extract from the adrenal cortex gland is used. Tablets of adrenal cortex in minute dosage are available for oral administration, but in the majority of cases suprarenal cortex liquid is preferable. The reason for this is twofold: (1) dosage is more easily regulated in the liquid form, and (2) 75 per cent of cases of this type also require insulin to inhibit an overactive thyroid and/or an overactive anterior pituitary.


Method to Maintain Blood Sugar at Normal Level—In cases of combined overactivity of the thyroid and anterior pituitary blood sugar levels may appear consistently normal. Insulin and suprarenal cortex dosage must then be so balanced in their ratio to each other that the blood sugar is maintained at normal. The expected

BIOCHEMICAL INDEX

BLACK LINE REPRESENTS PHOSPHORUS LEVEL — RED LINE CALCIUM LEVEL

DATE	B.S.	B.P.	U.S.G.	CA. P.	1	2	3	4	5	6	7	8	9	10	11	12
1st Blood																
11/5/48	100			10.4	3.7											10
2nd - In absence of sugar																
11/29/48	114			9.0	3.6											10

Length right forearm	Quotients	Length right leg
9.7	1.680	16.3
Wrist	A 1.475	9.3
one-fourth	B 1.180	7.15
one-half	C 1.229	9.4
three-fourth	D 1.274	11.2
Elbow	E	



Left forearm	Quotients	Length Left leg
Wrist	A'	
one-fourth	B'	
one-half	C'	
three-fourth	D'	
Elbow	E'	

•34•

Copyright 1950
By MELVIN E. PAGE, D.D.S.

reaction is for the cortex to raise the blood sugar and the insulin to lower it. A correct ratio of the two will maintain the blood sugar at the normal level. It is not surprising, therefore, to find an increase in the blood

sugar level in this case (Fig. 34) following the use of adrenal cortex alone.

Analysis of Figure 35

The outstanding physical characteristic of a person with a hypofunction-

ing anterior pituitary (DD' line) is the heaviness of the legs from the mid-point to the knee; there is no inward curve at the 3/4 circumferential measurement.


Results of Blood Analysis—In this

BIOCHEMICAL INDEX

BLACK LINE REPRESENTS PHOSPHORUS LEVEL — RED LINE CALCIUM LEVEL

DATE	B.S.	B.P.	U.S.G.	CA. P.	1	2	3	4	5	6	7	8	9	10	11	12
1st Blood																
7/19/49	100			11.4	4.2											14
2nd - In absence of sugar																
7/27/49	100			9.8	3.9											13

Length right forearm	Quotients	Length right leg
8.7	c 1.527	13.3
Wrist	A	
one-fourth	B	
one-half	C 1.427	13.9
three-fourth	D 1.490	15.5
Elbow	E	




Left forearm	Quotients	Length Left leg
Wrist	A'	
one-fourth	B'	
one-half	C'	
three-fourth	D'	
Elbow	E'	

•35•

Copyright 1950
By MELVIN E. PAGE, D.D.S.

BIOCHEMICAL INDEX

BLACK LINE REPRESENTS PHOSPHORUS LEVEL — RED LINE CALCIUM LEVEL



•36•

Copyright 1950
By MELVIN E. PAGE, D.D.S.

case (Fig. 35) the blood test prior to the use of anterior pituitary substance shows a blood sugar of 100, calcium 11.4, phosphorus 4.2, cell volume 40, sedimentation 14, blood pH 7.4.

Improvement Following Therapy—Anterior pituitary substance, 1/800 grain, was given daily for a week. Blood analysis made at this time shows a blood sugar 100, calcium 9.8, phosphorus 3.9, cell volume 40.9, sedimentation 13, blood pH 7.43. The .75 milligrams of excess calcium was reduced to nearly 0.

Endocrine Overdosage Should be Avoided—Since an overactive anterior pituitary, in the authors' experience, occurs in people with a susceptibility for the killing diseases (heart disease, cancer, diabetes) this agent, even in minute amounts, is used sparingly. The tendency in endocrine therapy should be to underestimate the dosage.

Analysis of Figure 36

The outstanding physical characteristic of a person with a hyperfunctioning anterior pituitary is an inward curve between the midcalf of the leg and the knee, at the 3/4 circumferential measurement. The difference in the appearance of the graph (Fig. 36)

at this point, compared with the previous graph (Fig. 35) will be readily discernible.

Results of Blood Test—Prior to administration of endocrine treatment, the blood test in this case showed a blood sugar 138, calcium 10.7, phosphorus 3.33, and cell volume 45.

Effects of Treatment—Daily use of 3 units of insulin for one week changed the blood picture quite radically: Blood sugar was 109, calcium 10.3, phosphorus 4, and the cell volume 45. Note that the blood sugar dropped 29 points and that the excess calcium was reduced 2 milligrams.

Blood Sugar Content Frequently High—This case illustrates the incidence of higher than normal blood sugar which is often encountered in people with hyperfunctioning anterior pituitary although medically they would not be classified as diabetic.

Insulin Used in Minute Amounts—Three to six units of insulin used to improve calcium-phosphorus blood levels in these cases seems to affect the blood sugar levels as well.

Effect of Insulin on Overactive Thyroid—In hyperfunctioning thyroid cases with normal blood sugar levels it is the authors' experience that the same minute amounts of insulin can

be given to inhibit the overactive thyroid and improve calcium-phosphorus blood levels without lowering the blood sugar levels.

In an article in JAMA³ it is stated: "Insulin, whether used in subcoma or in small tonic doses is a specific for the numerous every day nervous and physical complaints caused by habitual sympathetico-adrenal stimulation (prolonged alarm reaction)."

Analysis of Figure 37

A hyperfunctioning anterior pituitary (DD' line) may also respond to the use of hormones, as illustrated in Figure 37.

Blood Test—On the corrected diet the blood test shows blood sugar 100, calcium 9.6, phosphorus 2.38, hemoglobin 112, cell volume 47.7, blood pH 7.4, sedimentation rate 10.

Treatment—With one tablet of AD (an AD tablet contains what is estimated to be balanced amounts of both sex hormones in extremely minute quantities) the blood picture changed to the following: Blood sugar 100, calcium 9.8, phosphorus 2.66, hemoglobin 97, cell volume 42.8, blood pH 7.4, sedimentation rate 6. Excess

³Medical Literature Abstracts, JAMA 151:771 (Feb. 28) 1953.

calcium was reduced from 3.75 milligrams to 3 milligrams.

The pituitary is often called the "master gland" because of its influence on the growth and development of body tissues ("end-organs") and its stimulating power in "target glands" (thyroid, adrenals, gonads) through the action of tropic hormones.

As expressed by Wilkins:⁴ "The anterior pituitary gland acts as a control center of the system, and in turn is subject to regulation through the nervous system and by the concentration of circulating hormones elaborated by the target glands. It is apparent, therefore, that an alteration in one of the hormones affecting the body may be caused by a disorder arising in any of three centers of control: (1) the brain centers from which impulses pass to the pituitary, (2) the anterior pituitary, or (3) one of the "target glands"—thyroid, adrenal, or gonad. In diagnosis it should always be remembered that the primary lesion which gives rise to a particular hormonal disturbance may

be in any of these levels of control.

"Pituitary activities are regulated to a considerable degree by the blood concentrations of the hormones of the "target glands" [thyroid, adrenal, ovarian, testicular] and the pituitary may respond to some extent to alterations in the body chemistry which result from the excess or lack of these hormones."

Summary

The cases described illustrate the following therapeutic points:

(1) Thyroid extract augments thyroid activity; insulin inhibits thyroid activity. *Insulin is the opponent of thyroid.*

(2) Posterior pituitary substance augments posterior pituitary activity. *Adrenal cortex is the opponent of the posterior pituitary substance.*

(3) Anterior pituitary substance augments anterior pituitary activity; insulin, or the sex hormones, inhibits anterior pituitary activity. *Insulin and the sex hormones are the opponents of anterior pituitary substance.*

Abbreviated Statement of Formula

—If this formula is kept in mind the problems of treatment choice are minimized.

(1) *The thyroid is opposed by the insulin-producing cells in the pancreas.*

(2) *The posterior pituitary is opposed by the adrenal cortex.*

(3) *The anterior pituitary is opposed by either the insulin-producing cells in the pancreas or the sex hormones.*

Aids in Choice of Dosage—There are also several helpful clues in the matter of dosage:

(1) Degree of graph divergence from normal: the greater the divergence the larger the dose (all dosage is minute).

(2) Age: the younger the patient, the smaller the dose.

(3) With insulin correlate the degree of activity of the thyroid and/or anterior pituitary with the blood sugar level.


(4) With posterior pituitary extract correlate the degree of deficiency with the blood pressure and the specific gravity of the urine.

(5) With anterior pituitary substance correlate the degree of deficiency of the blood sugar. It may be low and with treatment it will rise; but *always err on the side of underestimating dosage of this endocrine.*

BIOCHEMICAL INDEX

BLACK LINE REPRESENTS PHOSPHORUS LEVEL — RED LINE CALCIUM LEVEL

DATE	B.S.	B.P.	U.S.G.	CA. P:	1	2	3	4	5	6	7	8	9	10	11	12
1st Blood																
5/26/53	100			9.6	238											
2nd - In absence of sugar																
6/2/53	100			9.8	266											



Objective of Technique—The purpose of this treatment approach is to use endocrines to supplement the glands' own production *only to the extent of their limitation*. A weak gland is much like a sick person, it needs good food (building material), it needs rest, and it needs its production load lifted. The following measures are intended to provide these benefits:

- (1) Correct the diet.
- (2) Include all the nutritional essentials for strength, growth, and maintenance.
- (3) Eliminate all things burdensome to glandular activity (primarily refined sugar).
- (4) Supplement the gland's own production by adding some of its own extract.

Necessity for Moderation Emphasized—To give endocrines in amounts in excess of need or in excess of normal total production would defeat the purpose of this treatment technique. The objective is to increase the efficiency of body chemistry. Too large a dosage of endocrines would create as great a problem in chemical adjustment for the mechanism as an inadequate production on the part of the gland itself.

On the matter of dosage Gordan and Lissner write:⁵ "Since hormonal agents are given to produce specific effects, it is usually possible to determine the proper dose for the particu-

⁵Gordon, Gilbert S., and Lissner, H.: *Endocrinology in Clinical Practice*, Chicago, The Year Book Publishers, Inc., 1953, pp. 26-27.

lar patient by a 'clinical bioassay.' The usual procedure is to start with the dose which, in the physician's experience, is the smallest effective amount of the agent."

Established Norms Indicators of Response—Unfortunately, too little is known about the amount of daily production of hormones. The handicap of insufficient knowledge in this aspect of treatment can only be overcome by observing with meticulous care the response to endocrine treatment. All the indexes of efficient chemistry the norms of which have already been established can be used as indicators of progress.

(End of Part Four)

2810 First Street North.

Medical Conditions Complicating Oral Surgery

BENJAMIN ELIASOPH, M.D.

Complications of Oral Surgery

The most important of the immediate emergencies encountered is shock or vasomotor collapse. A certain amount of psychic shock accompanies the mildest dental procedure and varies tremendously with each patient. The psychotic, the psychoneurotic, and the alcoholic patient may be cooperative and outwardly little disturbed but at other times almost impossible to treat.

Symptoms Noted—The cold, wet, clammy hands, the sudden pallor or flush, the general malaise, profuse perspiration, rapid pulse, nausea, faintness, or restlessness are symptoms often noted. Even in a mild form they interrupt the scheduled procedure and are time consuming and troublesome. The milder and more common episodes are usually psychic in origin, transitory, and readily amenable to treatment. Mild stimulants, such as whisky or aromatic spirits of ammonia, having the patient lie down, and words of encouragement are helpful.

This may occur in diabetic patients taking insulin, in rare cases of hyperinsulinism, and in some people who quickly metabolize their small

reserve of glycogen. The mechanisms of the reactions in the first two varieties are better known but the third needs some clarification.

Glycogen Depletion—This usually occurs in active, slightly nervous people almost always at the end of the working day after they have been rushing to finish any business to keep their appointment with the dentist. Sometimes they forego lunch or the afternoon snack that they are accustomed to having. Apprehension about the operation may play a role in depleting the blood sugar.

Blood Sugar Lowered After Sweet Drinks—It has been shown that some nervous people, particularly those complaining of fatigue, may have a drop in blood sugar after taking a sweet drink. In that case the island cells of the pancreas are overstimulated. Atropin and phenobarbital may inhibit the vagal stimuli and arrest this process.

Symptoms Prolonged and Severe—The symptoms of hypoglycemia may become as alarming as those of an attack of coronary thrombosis. With a real attack of coronary thrombosis, however, there should be some chest pain. This pain is usually precordial and radiates, as a rule to the left

shoulder and even down the arm. Radiations of pain to the neck and lower jaw, when present, are more likely to be found with organic disease in the heart or mediastinum.

Procedure Suspended—The presence of cyanosis, dyspnea, vomiting, and irregular pulse precludes further surgical operations for the time being and requires the help of a physician. Uncomplicated compensated cardiovalvular disease seldom gives rise to any difficulty. After an attack of acute rheumatic fever or chorea, it is advisable to wait as long as possible before completing surgery of any kind.

Patients with angina pectoris are fully aware of this condition and are usually prepared in advance and carry their medication with them.

Many oral surgeons follow the same procedure as the general surgeons in preparing their patients for operation:

1. They try to operate at a time when the patient is neither dehydrated and starved nor after he has taken a full meal.

2. Some make it a practice to give the patient an adequate dose of a barbiturate in order to achieve the

(Continued on page 568)

TERRAMYCIN

Used in Endodontic Cases

HARRY MAETH, D.D.S., Mosinee, Wisconsin

DIGEST

The history of the management of two root canal cases which were complicated with fulminating facial cellulitis of dental origin, followed by considerable suppuration, is described. In the past the author has occasionally extracted teeth in such condition, considering them hopeless for treatment.

In the cases reported in this article the antibiotic, terramycin, was employed as the exclusive therapeutic agent to control the acute inflammatory process, to sterilize the pulp canals, and as a topical postoperative dressing in the surgical wound area of root-end debridement.

Considerations in Endodontic Therapy

Endodontic therapy may be viewed as based upon radiographic, bacteriologic, and pathologic conclusions. To the specialist in endodontia, each of these considerations has been a link in the chain of his treatment.

Helpful in General Practice—To the general practitioner the radiograph offers information, although limited. Accurate bacteriologic determinations, however, and histopathologic studies are infrequently feasible or possible.

Many Possibilities in Treatment—The properties and limitations of cer-

tain therapeutic agents are well known to those practitioners with long experience in endodontic problems. Here again, however, the general practitioner is confronted with many theories and recommendations for treatment with chemotherapeutic agents; and more recently the antibiotics are available.

Case One

The patient in this case was a white female, age 18 years.

Symptoms—1. When first seen, August 1, 1952, swelling was present on the right side of the face, extending to the upper lip, cheek, and infraorbital area.

2. The temperature was 100.4° Fahrenheit.

Results of X-ray Examination—An extensive radiolucent area was revealed about the apex of the maxillary right central tooth, extending to the area of the root of the adjoining lateral incisor tooth. The lateral incisor responded normally to the pulp test while the central incisor did not.

Treatment—1. Terramycin, 500 milligrams, was administered orally as the initial dose, followed with 250 milligrams every five hours for the next three days.

2. August 4, 1952, dosage of terramycin orally was reduced to 250 milligrams every six hours.

3. Degenerated pulp of the maxillary right central incisor was re-

moved. The labial mucosa was incised at this site to establish drainage.

4. The pulp canal was irrigated with a solution of terramycin made from a tablet dissolved in distilled water. A cotton point saturated with the aqueous solution of terramycin was sealed in the root canal.

5. Terramycin cones were placed in the periapical incised area. A rubber dam drain was inserted.

6. August 6, 1952, treatment of terramycin was changed in the root canal. Terramycin medication was inserted into the surgical wound area.

7. On August 9, because of recurrent swelling at the labial aspect the incised area was opened wider for more adequate drainage.

8. Terramycin orally was continued, 250 milligrams every six hours. Drainage was maintained with a rubber dam drain.

9. August 12, 1952, the root canal was filled with a mixture of terramycin dissolved in beechwood creosote, combined with zinc oxide, and reinforced with fiberglass (hammermilled form). The necrotic soft tissue lesion was removed from the involved periapical area. The wound was sutured.

Histopathologic Report—The following report Number 12417 of the periapical tissue was made August 14, 1952 by the Laboratory of the Wisconsin State Board of Health, and signed by W. D. Stovall, M.D., and Gordon Worley, M.D.:

"Inflammatory tissue reaction. An epithelial lining of a cyst is not

found in the specimen submitted. No evidence of malignancy."

Case Two

In this case the patient was a white female, age 34 years.

Symptoms—When first seen, October 3, 1952, swelling on the right side of the face was present extending from the lip to the cheek and the infraorbital area. The temperature was 99.6° Fahrenheit. Considerable pain was experienced.

X-ray Examination—A large radiolucent area about the apex of the maxillary right lateral incisor, suggested a rarefying osteitis.

Treatment—1. Terramycin was administered orally, 500 milligrams followed with 250 milligrams every six hours. Swelling and pain subsided in less than twenty-four hours.

2. The pulp was removed October 3, 1952, and the root canal was treated with terramycin (soluble tablet) dissolved in sterile water. Terramycin was carried on a saturated cotton point to the root canal and sealed within.

3. Treatment to the root canal was changed the following day. Terramycin was continued orally, 250 milligrams every six hours.

4. October 7, 1952, treatment to the root canal was changed. Terramycin was continued orally, 250 milligrams every six hours. The same treatment was administered October 9 and October 11.

5. On November 11, 1952 after illness of the patient the treatment of October 11 was found intact. At this time the root canal was filled with a mixture of terramycin dissolved in beechwood creosote, combined with zinc oxide, and reinforced with fiber-

Case One

1. August 1, 1952, preoperative condition.

2. August 12, 1952, postoperative condition.

3. November 27, 1952, postoperative.


4. December 23, 1952, postoperative.

5. January 3, 1953, postoperative.

6. February 1, 1953, postoperative.

7. April 4, 1953, postoperative.





Case Two

8. October 3, 1952, preoperative condition.

9. October 11, 1952, during treatment. The canal is ready for filling.

10. November 1, 1952, the root canal filled.

11. February 8, 1953, postoperative.

12. April 3, 1953, postoperative.

3. No other therapeutic agent was applied during the entire course of treatment in both cases. (The beechwood creosote was used as the vehicle for the zinc oxide canal filling mass.)

Other Uses—Recent controlled hospital studies indicate the effectiveness of terramycin as the exclusive medication for the management of cellulitis of the face of dental origin.¹

Adjunct to Surgical Treatment—1. Terramycin has been investigated as an adjunct to surgical treatment where early diagnosis and prompt surgical intervention is made.²

2. Terramycin is effective against gram-positive and gram-negative organisms, and mixed infections.³

3. An increase in the incidence of penicillin-resistant strains of staphylococcus aureus has been observed during the past few years.⁴ This situation would suggest the advisability of using terramycin for topical and systemic therapy in a surgical infection, such as one which accompanies an involved dental pulp.

Prognosis

The prognosis of the two root therapy cases reported here is apparently hopeful. From a purely clinical standpoint it appears advisable to follow a course of therapy which assures definite progress.⁵

Mosinee, Wisconsin.

glas (hammermilled form). The necrotic tissue mass at the involved periapical site was extirpated and the wound was sutured.

Terramycin Successful in Therapy

1. It should be noted that after terramycin was administered orally

the acute stage in both cases subsided rapidly.

2. It is important to realize that terramycin probably inhibited and controlled the process of infection which was responsible for (1) the acute inflammation, (2) the rarefying osteitis, and (3) the degenerated pulp tissue.

¹Pollock, S. L., and Archer, W. H.: Terramycin in Dentistry, Pennsylvania State D. J. 20:No. 3 (March) 1953.


²Welch, H., and Lewis, C. N.: Antibiotic Therapy, Washington, D.C., The Arundel Press, Inc., 1951, chap. 15 p. 349.

³Welch, H., and Lewis, C. N.: Antibiotic Therapy, *ibid.*, p. 352.

⁴*Ibid.*

⁵Personal communication from William Ward Wainwright, D.D.S., University of Illinois, College of Dentistry, 1952.

1




Clinical and Laboratory

A Plaster Splint for Putrescent Pulp

Leo O. Beldo, D.D.S., Marquette, Michigan

1. When opening into the pulp chamber of a putrescent pulp the tooth is usually tender and painful. To immobilize the tooth, flow some fairly thick plaster on the labial surfaces including the teeth on each side of the one being operated upon. After the plaster has set, apply finger pressure on the labial of the splint and open into the pulp from the lingual.

2



Increasing the Rotating Speed of a Bur

I. B. Hyams, D.D.S., Montreal

2. The RPM of the bur may be increased by diminishing the circumference of the handpiece wheel. The speed will be doubled by reducing the circumference by one-half. Attach the wheel to a lathe on a universal chuck having previously removed the screw. Holding a rattail file against the revolving wheel, reduce the circumference of the wheel by one-half. Reduce the length of the screw and reassemble.

3



A Sure-grip Sprue

H. H. Look, D.D.S., Denver

3. A used Number 35 or Number 37 inverted cone bur is used as a sprue for wax patterns. The embedded inverted cone bur in the wax permits removal of the pattern without danger of the sprue and pattern separating.

READERS are Urged to Collect \$10.00

For every practical clinical or laboratory suggestion that is usable, DENTAL DIGEST will pay \$10.00 on publication.


You do not have to write an article. Furnish us with rough drawings or sketches, from which we will make suitable illustrations; write a brief description of the

SUGGESTIONS . . .

Patching an Alginate Impression

Harold S. Jones, D.D.S., Allentown, Pennsylvania

4. To correct an air bubble in an alginate impression, fill the void with any acrylic liner or self-curing restorative material by using the brush technique. The alginate impression should be dry before the correction is made.




4

A Simple Mount for X-rays

Daniel F. Tobin, D.D.S., New York

5. It is desirable at times to take x-rays and either project them by using a 2-by-2 projection machine or by placing them on a table enlarger. A convenient mount may be made by using two 2-by-2 pieces of thin cardboard stapled on three sides with the fourth side open. The windows are cut before assembling.




5

A Hanger for X-ray Films

S. Joseph Bregstein, D.D.S., Brooklyn, New York

6. A chrome tie rack with a swivel joint is used. This is useful for hanging x-ray films on the racks for drying. The tie rack can be attached to the wall in the processing room and when not in use it will lie flat against the wall.



6

technique involved; and jot down the advantages of the technique. This shouldn't take ten minutes of your time. Turn to page 558 for a convenient form to use.

Send your ideas to Clinical and Laboratory Suggestions Editor, DENTAL DIGEST, 708 Church Street, Evanston, Illinois.



Senile Anorexia

Patients showing the infirmities of old age usually have diffuse subjective complaints. The most common of these are: (1) a tendency to feel cold, (2) a distaste for living, (3) burning of the tongue, (4) a dry mouth, (5) loss of taste, and (6) difficulty of swallowing.

In many patients there is a notable lack of activity. Patients may spend their days lying in bed or sitting in a chair, apathetic, often without any initiative whatever, even with regard to eating or keeping clean.

The nutritional status varies. At times there may be remnants of previous obesity, but more often the patient is emaciated, more rarely cachectic. The skin is pale, more so than can be accounted for by the hemoglobin percentage which usually ranges between 70 to 85 per cent. X-ray films of the vertebral column often reveal osteomalacia. Chronic diarrhea and recurrent pyuria are common complications.


The course is typically chronic. The condition may last for years but the stupor usually increases and the patients develop decubital ulcers and incontinence of urine and feces. Prognosis is poor and patients seldom regain their strength if left untreated.

The debility is associated with a group of symptoms not found in active persons, even if they have attained an extremely advanced age. These symptoms are those noted in glossitis; for example, (1) marked reddening and atrophy, (2) hypertrophy of the papillae of the tongue, (3) cheilitis and bright red lips, (4) desquamation of the skin in the angles of the mouth, and (5) perleche.

Symptoms of nutritive failure are common among extremely old persons living by themselves. The daily food consumption is definitely deficient. It is believed that senile debility may be caused by a primary disturbance of the appetite regulation associated with various medical and

MEDICINE

and the Biologic Sciences



psychologic factors which are peculiar to old age. These factors tend to reduce both the quantity and quality of the food consumed.

Vinther-Paulsen, N.: Senile Anorexia, Geriatrics 7:274-279 (September-October) 1952.



Sciatica

The term is used to designate pain which begins in the back; namely, the sacroiliac region. The pain extends into the buttocks, the posterior or posterolateral aspect of the thigh at least as far as the knee, often into the posterior or posterolateral aspect of the calf and ankle (either or both) and even out into the foot. The disorder is quite common.

Anatomically the cause lies in one of three places: (1) in the sciatic nerve, (2) in the lumbosacral plexus, or (3) in the nerve roots that ultimately make up the sciatic nerve; the fourth and fifth lumbar, and the first sacral nerve roots. The symptoms of

80 per cent or more of all patients who have sciatica are attributable to nerve root lesions.

The nerve root lesion is sometimes referred to as "radiculitis." This implies that one of the radicals or nerve roots is being irritated by some pathologic process. A number of disease entities can produce a nerve root lesion. The most common of these is a protruded intervertebral disc. About 90 per cent of all lumbar and upper sacral nerve root lesions are attributable to protruded intervertebral discs.

Other disorders that can cause such lesions are (1) spondylolisthesis, (2) tuberculosis with Pott's disease, (3) osteomyelitis, (4) primary and metastatic neoplasm, (5) epidural abscess, and (6) diabetes.

Usually there are two components to the pain: 1. More or less constant aching, burning discomfort with stiffness low in the back continuing along the course of the sciatic nerve. The pain extends to the knee. 2. Increased severity of the pain which extends or travels inferiorly into the thigh, past the knee into the calf, and sometimes into the foot and toe. This shooting pain is almost always present during an exacerbation of the disorder.

Treatment depends on the etiology. A neoplasm in the thigh pressing on the sciatic nerve should be removed. If there is a metabolic dysfunction it should be corrected. Diabetes should be controlled. Vasodilators should be given in case there appear to be some vasospastic phenomena. When malignancy attacks the lumbosacral plexus in the pelvis about all that can be done is in the form of palliation.

The protruded intervertebral disc must either be left in or taken out. About three-fourths of the patients receive a satisfactory result from surgery. Few patients who have protruded intervertebral discs ever ultimately have a completely normal back which will stand any kind of punishment. This fact should be realized early by each patient.

Millikan, Clark H.: Causes and Treatment of Sciatica, Postgrad. Med. 12:207-211 (September) 1952.



Lobotomy— Present Status

Prefrontal lobotomy is effective therapy in severe cases of schizophrenia, especially when the patient is obstreperous, anorexic, and restless. Persons with the following symptoms: (1) anxiety neuroses, (2) obsessive tension states, (3) depersonalization syndromes, (4) agitated depressions, and (5) chronic pain with or without organic basis, obtain relief from the operation.

An active and conscientious lobotomy program can transform a mental institution into an old people's home.

Hyperactive subjects show remarkable reduction in disturbed and disturbing behavior after lobotomy. These patients may not be able to adjust to conditions outside the hospital. Their relations with other patients, however, are much improved and simple duties can be performed. The operation often brings relief of suffering, loss of fear, and therefore loss of hate.

The patient who is able to live at home or even work may not gain from the operation. He may become idle, outspoken, tactless, and irresponsible. Lobotomy is not advisable for these better preserved patients.

The operation must be extensive enough to control the symptoms. If it is too extensive it is ruinous. Conservative operations are better than extensive ones in the following situations: (1) elderly people, (2) severe painful conditions, (3) psychoneurosis, and (4) agitated depression.

Lobotomy should be avoided in cases of (1) alcoholism, (2) drug addiction, except in pain conditions, (3) criminal psychopathy, and (4) organic brain disease except parkinsonian and thalamic syndromes.

Primarily, lobotomy will reduce painful effect and eliminate dread and tortured self-concern. The more anxiety the patient had before the operation, the more dramatic the improvement. Depression, with a fear of the future based on fancied past misdeeds, yields magically.

Most patients with (1) delusions, (2) phobias, (3) conversion reactions, (4) obsessions, and (5) many psychosomatic complaints are helped. Those with hallucinations may benefit early in the disease, but effectiveness of the procedure lessens as the symptom continues.

Cases of early schizophrenia do well with lobotomy. However, many of these patients are seen only after weeks or months of other therapy including electroshock, which may cloud the symptoms and conceal the advance of deterioration.

A six-month's trial of conservative care in an institution should be assayed before considering surgery. Decision should be made soon thereafter. After a year in the hospital the chances for recovery rapidly diminish.

Freeman, Walter: *Present Status of Lobotomy*, Virginia M. Monthly 79:436-439 (August) 1952.



Tetanus

Tetanus is one of the greatest hazards of wounds. It is sound practice to assume that any open wound resulting from trauma is vulnerable to tetanus infection. Such a wound should be an indication for a booster dose of toxoid or prophylactic antitoxin.

In the United States Army active immunization with tetanus toxoid is considered effective for four years. Booster doses given at the time of injury provide a prompt increase in circulating antibodies.

The alum-precipitated toxoid is slowly released from the site of deposition. Immunization is completed with two injections at intervals of from four to six weeks. The fluid toxoid is more rapidly absorbed, is less apt to produce cold abscesses, and is given as a series of three injections at intervals of four weeks.

Passive immunization with tetanus antitoxin is less certain and less effective and lasting than active im-

munization. Use of tetanus antitoxin is usually limited to two situations: (1) Treatment for clinical tetanus, and (2) occurrence of wounds or burns necessitating protection against tetanus in patients who have not been previously immunized or who have not completed initial active tetanus immunization.

The prophylactic dose for adults is 3,000 units in treatment for clinical tetanus. If the patient is treated one or two days after injury, the prophylactic dose should be increased to 10,000 to 20,000 units. The duration of passive immunity induced thereby is five to ten days. If the wound is not clean by this time, further passive immunization is indicated.

When a diagnosis of tetanus has been determined and preliminary testing for sensitivity to horse serum has been done, tetanus antitoxin is given. Fifty thousand to 100,000 units intramuscularly constitute a dose large enough to neutralize all the unfixed or free toxin in any patient who has a chance for recovery. From 10,000 to 20,000 units intramuscularly are administered for the following four or five days also. Appropriate antibiotic therapy is also indicated.

Pulaski, Edwin J.: *Antiseptics, Antitoxins and Antibiotics*, Mod. Med. 20:84-90 (September 1) 1952.

This is a Human Body

IN 1928 STUDIES were being made from the point of view that a human being is a product of the air he breathes, the water he drinks, and the food he eats; day after day, year after year, the tissues of the human body are replaced by intricate combinations of oxygen, water, and food. Today this is a medical axiom, and physicians have gone a long way toward showing that disease is chemical and chemically correctable.

—Tom D. Spies, M.D.: JAMA 153:185-193 (Sept. 19) 1953.

The EDITOR'S Page

"IN THIS age of addition, if anything is wrong with a patient that is not fully understood the tendency is to seek the cause not among the harmful things that may be present but among the 'good' things that may be absent. And so one supplement after another is added. The scientific possibility of *subtraction* apparently has occurred to no one."

This quotation is from the important work of Charles Dillon, D.D.S., L.D.S.¹ of Scotland and is a profound observation of present trends in therapy. Let us look around us and see how accurate this observation is. To mind come the additives: vitamins for deficiencies in food, antibiotics to counteract bacterial activity, the fluorides to reduce dental caries.

It is easier and more lucrative to prescribe the addition of something than to advise the subtraction of something. For example, in the field of nutrition, it is simpler for the dentist and the physician to suggest the ingestion of the synthetic vitamin products than it is to advise a complete change in dietary habits by the removal of noxious foodstuffs such as refined sugar and white flour. Entirely within the field of economics it is more acceptable to the patient to be given a positive suggestion and a prescription to *take* something than it is to be told to refrain from acts of eating that are pleasurable. People are more disposed to take a pill or any other form of medication and believe in its magic than they are to discipline themselves to forego an experience.

The antibiotics, boons to mankind that they are, have blunted the zeal for accurate diagnosis. When a person complains of malaise, or pains, or disturbances of any function he is given an antibiotic. When the body temperature is elevated slightly, or when the white blood cell count is increased, he is given an antibiotic. When gums are inflamed or bleeding, when irritation appears in the residual dental ridges, the patient is given an antibiotic. Each one of these procedures represents sloppy diagnosis and "shotgun" therapy.

Fluoridation is certainly an example of the addition of a chemical to control a dental disease the nature of which we do not fully understand. The presumption is that by some process not altogether clear in our minds the structural resistance of the tooth can be improved to better withstand the assaults of bacteria on the carbohydrates that produce lactic acid. Although we know that the caries rate may be reduced by the *subtraction* of the sugar available for degradation and acid formation, we also know that this discipline is not readily accepted by patients. People would much prefer to receive *additive* chemicals in painless doses in their drinking water and continue to use sugar without *subtractive* limitations.

It is regrettable that the subject of fluoridation has lost scientific objectivity and that so much emotion has been engendered. This is not a subject to be treated with passion. Instead it requires scientific detachment. Not so much "pro" and "anti" as "how" and "why."

We have not made clear enough to the public that *reduction* in the caries incidence is different from *prevention* of caries. We must prepare the public for this fact. Dillon uses these words to describe this:

"If lactic-acid fermentation is going on actively in the oral cavity it matters not whether the teeth are well calcified or poorly calcified, or whether 'hardened' by fluorides; it is only a question of time and the rate of acid fermentation, before the inorganic superstructure is breached, permitting the microbes to enter the basic organic structure, and from then on the process is one of progressive protein hydrolization, and the only way to inhibit it is to find some safe way of controlling fermentation."

Fluoridation is not a panacea or a total preventive. At best it is a valuable agent to reduce the caries rate; at the worst, it is a cumulative toxic agent. Time will expose the answer.

¹Dillon, Charles: Fluorine and Dental Caries, Dental Practitioner 3: No. 3, 79-86 (Nov.) 1952.

Annual Index-1953

Anatomic Reconstruction

- Goldman, Irving: The Art and Science of Full Mouth Rehabilitation, January 17
Shohet, Harmon: Fundamentals in Mouth Reconstruction and the Temporomandibular Joint Syndrome—Part One, June 257
Shohet, Harmon: Fundamentals in Mouth Reconstruction and the Temporomandibular Joint Syndrome—Part Two, July 307

Anesthesia

- Brudzynski, C.: Investigation of Anesthetic Deaths (An Abstract), March 110
Faier, A. D.: Supplemental Local Anesthesia, June 248
Millhon, Jerry A.: Pentothal Sodium Anesthesia for Dental Surgery in Office Practice (An Abstract), June 264
Policow, Myron: Analgesia in Dentistry, September 391
Shaw, J. C. Middleton, and Dodds, A. E.: Hypotensive Anesthesia in Dentistry (An Abstract), August 347

Caries

- Dental Caries in Rats Fed a Diet Containing Processed Cereal Foods and a Low Content of Refined Sugar (An Abstract), August 360
Mechanism of Cariostatic Action (An Abstract), January 27
pH of Saliva and Tooth Decay (An Abstract), August 366

Clinical and Laboratory Suggestions

- January 28
February 76
March 122
April 174
May 222
June 268
July 316
August 358
September 402
October 456
November 504
December 548

Contra-Angles

- January
Profile of a Scientist; Sleep and What You Make of it 42
February
"A Negative Appetite for Activity"; The Line Forms on the Left 89
March
Hold That Advice!; Fate and Fluorides 136
April
Just Peanuts!; Personality Rehabilitation 184
May
Five Things that Physicians Shouldn't do to Dentists 232
June
Our Health and the Atomic Bomb; The Enemy, Sugar 281
July
Wisdom of India; Chicken Feed; Recent History 328
August
The Search for the Soul 368
September
Getting Mad is Bad 418
October
Two-thirds of the Young with Gum Disease; The One-string Cello Player 462

- November
Healthy Human Nature; My Friends David and Anne 510
December
Two Views on Nutrition 556

Dentures, Full and Partial

- Arrow, William: Laboratory Aspects in the Construction of Implant Dentures, June 254
Craddock, F. W.: The Forms of Porcelain Posterior Teeth, February 62
Implant Dentures May be Vindicated (An Abstract), July 327
Lew, Isaih, and Kestenbaum, Irving: An Implant Button Technique for Denture Prosthesis, July 298
Moran, Hugh G.: Jaw Relations: Their Significance and Importance in Denture Construction, March 104
Myerson, Simon: Occlusal Forms and Nature's Plan, February 65
Page, Harry L.: Natural Articulation Movements, May 202
Page, Harry L.: A Critical Appraisal of Centric Relation, August 342
Smokler, Jack: Lingual Metal Casting, November 500
Spoon Dentures (An Abstract), July 306

Diagnosis

- Castigliano, S. Gordon: A Decalogue on Cancer for the Dentist (An Abstract), January 39
The Oral Cavity (An Abstract), September 390
Weinberger, Alexander: The Diagnostic Significance of Nutrient Canals, July 301

Editorials

- January
Fluoridation to Reduce Caries Incidence 30
February
The Human Uses of Chlorophyll Products 75
March
The Clinical Impression a Principal Source of Knowledge 124
April
The Theory of Focal Infection 176
May
The Dental Seminar 224
June
Anxiety in the Dental Patient 270
July
Reactions to the Use of Antibiotics 318
August
Degree of Efficiency in Denture Wearers 357
September
The Book of Health 404
October
The Effects of Electrolytic Action in Dental Restorations 453
November
Hazards to Dentists from X-ray Irradiation 503
December
Changing Implications in Treatment 552

Endodontics

- Maeth, Harry: Terramycin Used in Endodontic Cases, December 545

Medical Subjects


- An Evaluation of Calcium and Phosphorus Medication in Dentistry (An Abstract), January 40
Chemistry of Disease (An Abstract), December 535

- Dillon, Charles: The Biochemistry of Fluorides, November 486
Dillon, Charles: Detoxifying Function of Bone (An Abstract), November 489
Dillon, Charles: Retention of Fluorine (An Abstract), November 489
Dillon, Charles: Mottling (An Abstract), November 490
Erythrocyte Sedimentation Rate (E.S.R.) (An Abstract), November 520
Fluoridation of Water Supplies (An Abstract), April 161
Headaches and Facial Pains in Lesions of Cervical Discs (An Abstract), March 134
Johnson, W. Basil: Antihistamine Drugs to Inhibit Edema Following Oral Surgery, July 314
Kinawi, M. M.: Ethyl Chloride Spray in the Treatment of Fibrositis, Myositis and Allied Conditions (An Abstract), August 356
Laszlo, Daniel, and Spencer, Herta: Medical Problems in Management of Cancer (An Abstract), November 516
Page, Melvin E., and Brooks, D. L.: Body Chemistry in Health and Disease—Part One, September 394
Page, Melvin E., and Brooks, D. L.: Body Chemistry in Health and Disease—Part Two, October 440
Page, Melvin E., and Brooks, D. L.: Body Chemistry in Health and Disease—Part Three, November 491
Page, Melvin E., and Brooks, D. L.: Body Chemistry in Health and Disease—Part Four, December 536
Smith, W. Hamilton: Mephenesin® in the Treatment of Tetanus (An Abstract), September 416
Strean, Lyon P., and Horton, Charles Pinckney: Hydrocortisone in Dental Practice, January 8
Strean, Lyon P.: Vitamin B₁₂ in Tic Douloureux, May 214
This is a Human Body (An Abstract), December 551
Water Fluoridation: The Respective Responsibility of the Health Profession and the Community (An Abstract), May 213

Medicine and the Biologic Sciences

- January
Nail Shapes; Transmission of Poliomyelitis Virus; Iron Metabolism; Hypothyroidism in the Aged; Reading Difficulty; Gout—Recent Advances; Surgical Treatment of Pain 31
February
Basal Metabolism Test; Blood Sedimentation Rate—Importance; Boric Acid Poisoning; Venous Thrombosis; Urinary Calculi; Local Anesthetics—Toxic Reactions; Fetal Deaths; Syphilis Treatment; Pregnancy After Forty 78
March
Status of Vagotomy; Sunglasses; Anemia in Pregnancy; Needle Biopsy of the Liver; New Prescription Regulations; Varicose Veins; Reconstructive Surgery for the Elderly; Carcinoma of the Lung—Incidence 126
April
New Tuberculosis Cure; The Diabetic Diet; Irradiation Dermatitis of the Hands; Pain in the Aged; Sigmoidoscopy—Value; Bone Grafts 177

KODACHROME SLIDES available



Sixteen of the charts in the series *Visual Education in Dentistry* are available in KODACHROME slide form. The size is the usual 2" x 2" and the slides fit any standard projector.

The titles of the slides are given in the box to the right. The slides are reproductions of sixteen of the most popular charts in *Visual Education in Dentistry*. Many dentists have been using the complete booklet in their practices for quite a few years and total sales now are approximately 110,000.

You will find the KODACHROME slides most effective for use at the chair, in explaining dental conditions to patients. The series is also extremely valuable to dental societies, dental instructors, and clinicians.

The price is only \$16 and this includes appropriate lecture material. The coupon is for your convenience.

Titles of Slides

1. Dental Conditions
2. The Progress of Tooth Decay
3. Pyorrhea Treated or Neglected
4. What Does An X-Ray Show?
5. "Things Are Not Always What They Seem . . ."
6. The Requirements of a Correct Restoration
7. Insulation
8. How Irregularities of the Teeth Affect the Face
9. How the Loss of Teeth Affects the Face
10. The Collapsed Face
11. The Action of Local Anesthetics
12. Why Construct a Bridge?
13. The Danger From the Impacted Tooth
14. The Circulation of the Blood
15. "One Rotten Apple Can Spoil a Bushel"
16. "A Little Neglect May Breed Mischief"

Dental Digest, 1005 Liberty Ave., Pittsburgh 22, Pa. D.D.12

- ☐ Here is my check in amount of \$16. Please send in 16 KODACHROME slides and lecture material promptly.
- ☐ Bill my dealer.

Dr. _____

Address _____

City _____

State _____

Dealer _____

May	Heart Failure and Environmental Heat; Glaucoma; Obesity; Constipation; Chronic Edema of the Lower Limbs	226
June	Deterioration of the Bedfast Patient; Influence of Television on Children; Male Fertility; Vascular Disease in Diabetes; Thin Bones; The Thyroid and the Nose	274
July	Effects of Tobacco Smoking; Antibiotics—Effective Use; Poisoning in Children; Cerebral Palsy—Etiology; Mental Health and Music; Epilepsy	321
August	Restriction in Cardiac Cases; Antisepsis in Surgery; Antibiotics for Prophylactic Use; Radiation Necrosis of Jaw Bones; BCG Vaccination;	

Backache; Mirth Response Test	361
September	
Peptic Ulcer—Surgical Treatment; Pain-relieving Drugs; Poliomyelitis and Injections; Multiple Sclerosis—General Consideration	401
October	
Tinnitus Aurium; Curare in Poliomyelitis; Stimulation of Growth in Long Bones	458
November	
Intraoral Cancer—General Considerations; Maxillary Fractures; Antibuse® Therapy	506
December	
Senile Anorexia; Sciatica; Lobotomy—Present Status; Tetanus	550
Miscellaneous	
Announcement of Books Received, June	253

Announcement of New Publication, Excerpta Medica, Cancer Section, October	4-5
Annual Index: December	5-3
Civil Malpractice (An Abstract), December	5-5
Cleft Lip and Palate and Parental Age (An Abstract), May	2-21
Difference in Obligations of General Practitioner and Specialist (An Abstract), November	4-90
Endocrine System (An Abstract), November	5-15
Gamma Globulin (An Abstract), August	3-53
Green Teeth (An Abstract), March	1-36
King, Charles Glen: Longevity and Nutrition (An Abstract), November	4-97
Mellanby, Sir Edward: Chemical Manipulation of Food (An Abstract), June	2-71
Moore, W. Amos: Rotated Bicuspid and Axial Gradients, August	3-48
Related Syndromes (An Abstract), September	3-92
Seltzer, Samuel: Tray For Use in the Practice of Endodontics, June	2-65
Sheppard, Irving M.: Tongue Dynamics, March	1-117
Superior Visceral Skeleton (An Abstract), April	1-159
Syphilis as an Occupational Hazard to Physicians (An Abstract), June	2-67
Tonsillectomy and Poliomyelitis (An Abstract), August	3-67

Operative Dentistry

Coggeshall, R. J.: Costs and Other Considerations in Operative Airbrasive® Technique, April	1-68
Feldman, Francis, and Kay, Samuel: Cast Splint Retained by Horizontal Pins, May	2-18
Hamburg, Harold L.: Fixed Bridge Assemblage, April	1-60
Miller, Charles Jay: The Tube Impression, April	1-171
Physiologic and Neurologic Occlusion (An Abstract), December	5-62
Rubinstein, Murray N.: Mouth Rehabilitation in Practice, May	2-207
Silverman, Meyer M.: Gingival Pits for Better Amalgams, August	3-54

Oral Pathology

Fischer, Guido: Apical Region and Pulpless Teeth (An Abstract), July	3-19
Payne, Reginald T.: A Case of Lipoma of the Parotid Gland (An Abstract), September	4-12
Prevention of Cancer (An Abstract), May	2-220
Protzel, Martin S.: Herpes Zoster, August	3-49
Tam, Joseph C.: Lip Hemangioma Treated with Five Per Cent Sodium Morrhuate, August	3-55
Bruce, Karl W.: Mandibular Cyst of Uncertain Origin, November	4-98
Cardiac Patient (An Abstract), December	5-65

Oral Surgery

Chernoff, Phillip M.: Surgery vs. Bacteriology in Pulpless Tooth Management—Part One, January	2-23
Chernoff, Phillip M.: Surgery vs. Bacteriology in Pulpless Tooth Management—Part Two, February	2-71
Chernoff, Phillip M.: Surgery vs. Bacteriology in Pulpless Tooth Management—Part Three, March	3-111
Chernoff, Phillip M.: Surgery vs. Bacteriology in Pulpless Tooth Management—Part Four, April	4-162



This Ney-Oro G-3 spoke casting
demonstrates the great
margin of safety in
dentistry's finest partial
denture gold.

Ney-Oro A-1
for Inlays

Ney-Oro B-2
for Bridges

Ney-Oro G-3
for Partials



Control infection quickly with PHYLORINOL

An antiseptic solution powerful enough to destroy bacteria very quickly, yet so safe that it can be used full strength on delicate orificial tissue. Phylorinol is used in more and more dental schools and hospitals and is acclaimed by thousands of practitioners as the medication of choice as an adjunct in the treatment of

Gingivitis
Inflamed or Bleeding Gums
Vincent's Infection
Periodontal Pockets
Acute Third Molar Flaps
Malodorous Oral Conditions
Extractions
Dry Sockets
Surgical Care
Denture Soreness
Tooth Sensitivity
Caused by Receding Gums

The combination of iodophenol and chlorophyl is many times more effective than either ingredient alone. We would like to have you prove the efficacy of Phylorinol in your own practice. Simply send \$2.00 for a trial 8 oz. bottle. Your money

will be refunded if you are not entirely satisfied. No special instruments or skills are required. Non-toxic, non-irritating. This offer is made only to dentists and physicians. Phylorinol is never advertised to the laity.

SCHAFFER LABORATORIES, 3512 Ocean View Blvd., Glendale 8, California
I enclose \$2.00 for a trial bottle of Phylorinol on money-back guarantee of satisfaction.


Dr. Address
City State
My Dental Dealer is
D-15

PHYLORINOL

(Fill-or-inol)

SCHAFFER LABORATORIES

3512 OCEAN VIEW BLVD., GLENDALE 8, CALIFORNIA



Contra- Angles



Two Views on Nutrition

An example of divergent basic philosophies is expressed in two points of view on nutrition, one by an American scientist, the other in a discussion by members of the British House of Lords. The American is the scientific director of the Nutrition Foundation which is a group organized and richly subsidized by some of the largest manufacturers and processors of food. This point of view emphasizes nutritional surveys, research grants, and technologic developments to improve nutrition. The talks in the House of Lords were on a less exalted scientific level and were concerned with the potential dangers in processed and chemically treated foods. Here the desire expressed was: "To increase the availability of fresh foods for the people."

The American point of view, as stated by Charles Glen King, is in part as follows:

"From a food supply point of view, we in the United States are exceedingly fortunate. We have available enough of all the nutrients to furnish an ideal diet for the entire population, plus a reserve quantity for export. Nearly everyone has access to a 'free' education. Economic resources and transportation facilities are adequate to reach nearly everyone. Protection of food supplies by industry and by public law has set a high standard for several decades; and the foods available in our markets are as attractive and flavorsome as one could find anywhere. Yet we face such records as these:

"1. Apparently only about one-fourth of our population consumes a diet that well-trained scientists regard as good or excellent.

"2. At the opposite end of the picture, we have about 20 million or

- | | |
|--|-----|
| Eliasoph, Benjamin: Medical Conditions Complicating Oral Surgery (An Abstract), December | 544 |
| Holland, Daniel J.: Excision of Rantula, August | 352 |
| Ingle, John I., and Dow, Pierre: Replantation Following Total Luxation, September | 386 |
| Ogus, William I.: Unilateral Operation for Correction of Prognathism on Rotated Mandible, July | 304 |
| Paullus, W. S., and Gordon, J. T.: Surgical Procedure for the Lower Implant Denture, February | 56 |
| Samaha, Francis J.: Hemorrhage in Relation to Dentistry, April | 152 |
| Shaner, M. H.: Simplified Technique for Removal of Impacted Lower Third Molar, August | 346 |
| Sherman, J. H.: Foreign Body in Root Canal: Report of a Case, September | 393 |
| Orthodontics | |
| Klein, Earnest T.: Pressure Habits, Etiologic Factors in Malocclusion (An Abstract), April | 167 |
| Periodontics | |
| Bands On Teeth (An Abstract), January | 16 |
| Roentgenography | |
| Shapiro, I.: Film and Mechanical Aids in Intraoral Roentgenography, December | 530 |
| Waldman, David: X-ray All Delayed Eruption Cases, March | 114 |

more people who shorten their lives, handicap their health and bring on many of our most destructive diseases by consistently eating and drinking too much—that is, they consume too many calories. It is socially, medically, mathematically and anatomically correct to insist that these figures should be reduced!

"3. The incidence of tooth decay, most of which could be prevented by good nutrition, is extremely high. Although conditions are at last improving, most communities show an incidence of tooth decay (i.e., a record of one or more decayed, filled or missing teeth) in the range of 75 to 95 per cent of the population by the time adolescence is reached.

"4. The infant and maternal death rates are excellent compared with other countries, but most medical authorities agree that the record could be improved markedly by good nutrition. (We should have more counties reaching the Westchester County, New York, record of last year—reported as 'not a single instance of maternal death' in a county whose population is well over 600,000.)

"5. The top four causes of death in the United States—heart disease, cancer, cerebral hemorrhage and diseases of the new-born, and many of the diseases below fifth place (influenza and pneumonia) are metabolic, rather than infectious in nature. The instances of hypertension and gastric ulcers at meetings of Corporation Boards of Directors are too frequent for our comfort and economic advantage. These diseases are, on an experimental basis at least, subject to partial prevention and favorable therapy by means of good diets.

"6. Although we have not had recent adequate surveys of the nutritional status of our population, such data as we have, support the conclusions based on food habits, that low intakes of protective foods which furnish liberal quantities of vitamins, minerals and good quality proteins, are inexcusably common. The public, largely through ignorance and negligence, does not make intelligent use of the excellent food that is available. Our three worst enemies in the field

Amco-Facts

AMCO takes this opportunity

to wish all of you

A Merry, Merry Christmas

A Prosperous New Year

and

An Invitation

to investigate

Micro-seal

FOR FULL IMPRESSIONS


Our success, locally, has encouraged us to recommend Micro-Seal to you through this excellent journal.

Micro-Seal, an elastic impression material for full dentures, uses no water—thus has no dimensional change. You can pour your model days later, if necessary, and make more than one model! Micro-Seal may be used with positive assurance of accuracy.

An organosol with absolutely no burn or sting, Micro-Seal sets in the mouth in 2 minutes. There is no similar product on the market.

The versatility of Micro-Seal makes it adaptable to any technic.

You may order from your dealer with a cheerful "use or return" guarantee or we will send a generous sample for \$1.00.



AMERICAN CONSOLIDATED DENTAL CO.

809-11 N. NINETEENTH ST., PHILADELPHIA, PA.

12-53

CLINICAL AND LABORATORY SUGGESTIONS

(See pages 548 and 549)

Form to be Used by Contributors
To: Clinical and Laboratory Sug-
gestions Editor

DENTAL DIGEST
708 Church Street
Evanston, Illinois

From: _____

Subject: _____

Explanation of Procedure:

Sketch:

Suggestions submitted cannot be
acknowledged or returned.

\$10 will be paid on publication for
each suggestion that is used.



The last word in COPPER DIE & MODEL PERFECTION . . .

THE ORIGINAL
Silverizer



SUPERIOR Impression HOLDERS

With These Outstanding Improve-
ments:
Will not distort impressions during
mounting or dismounting operations.
Impressions can be mounted or dis-
mounted in an instant.
Light, simple and easy to handle.
Fits on all Electro-Formers.
Automatically holds impression and
makes visible electrical contact with
it.
Mounts impressions with or without
copper bands.
Made of special alloy, non-corrosive-
stainless steel. It has exceptionally
strong plastic insulation cemented on
with specially developed cement.
Made to last for a lifetime.
It comes with improved, new, time-
saving technique directions.

Only \$2.25 each
plus postage

Be ware of higher priced crude imitations that undersize the dies.
Buy Impression Holders and Silverizer. We guarantee you
satisfaction or money refunded.

SUPERIOR ELECTRO-DENTAL & MEDICAL APPLIANCE CO.
307 WEST FIFTH STREET BEAUMONT, CALIFORNIA
Since 1936 Manufacturing Superior Copper Die Electro-Formers in 4 Models,
For 6, 12, 24 and 30 Dies.

One patent allowed; other patents pending
FOR METALIZING DENTAL IMPRESSIONS, FOR
ELECTRO-FORMING MICROSCOPICALLY ACCUR-
ATE COPPER DIES AND MODELS.

GUARANTEED BEST!

Silverizer Powder & Thinner Liquid, Tech.
B is perfected for use in the customary two
step application Technic.

Silverizer Fluid, Tech. C is perfected for
use in a simple one step application Technic.
SILVERIZER IS LOW PRICED, VERY
ECONOMICAL IN USE, AND IT PLATES
BETTER WITHOUT TROUBLES.

It took us many years to
perfect Silverizer.

of nutrition are first, complacency;
second, delays in developing adequate
programs of basic and applied re-
search; and third, failure to develop
adequate education.

"Practical goals that we could
reach within the next ten years are of
major importance in the health, econ-
omy, and total strength of the nation.
We could accomplish:

"(1) A sharp decrease in the death
rate from heart disease, hardening of
the arteries, high blood pressure,
cerebral hemorrhage, diabetes, and

liver disease, plus a lower incidence
of cancer. This result would be re-
flected in an extension of the years of
vigorous living—by deferred aging—
not simply by added years of survi-
val.

"(2) A steady lowering in the ma-
ternal and infant death rate. The re-
productive cycle represents a normal
physiologic stage of life, but it does
impose a requirement for a higher
level of nutrition than is required of
adults at other times.

"(3) The incidence of tooth decay

Hygienic COLD CURE Denture Acrylic


This fine new self-curing Denture Acrylic is ready to finish
after 60 minutes in the press. No heat required. The finished
denture is hard, dense and takes a beautiful polish. Same shades
and granulations as regular Hygienic (Hot Cure) Acrylic and
may be used interchangeably.

Try this time and labor saver. Include some in your next order
from your local dental dealer.

For Free Sample write —

The Hygienic Dental Mfg. Co.
Akron 8, Ohio, U. S. A.

DD-1253



should drop to only five or ten per cent of the present level. About half the goal can be reached by fluoridation of water supplies, the balance chiefly by improved total nutrition.


"(4) The day-to-day level of mental and physical performance of our population would rise sharply by avoiding the transitory periods of low efficiency that result from poor food practices, such as omitting breakfast or lunch, and stuffing at other meals.

"(5) Thyroid gland enlargements or goiter caused by low iodine intake should completely disappear, along with scurvy, rickets, pellagra, nutritional anemia, subclinical beriberi and other diseases caused by specific deficiencies. Iodized salt, citrus fruits and juices, vitamin D milk, green leafy and yellow vegetables, enriched cereals, and meat are major factors in accomplishing these advances.

"(6) The diseases associated with glandular failure, such as arthritis and many nervous disorders, should decrease markedly as a direct result of balanced diets. In this area of public health, we are in the most poorly defined of all the fields mentioned. These gains are available at levels of nutrition well above the intakes that cause so-called deficiency diseases. Their continued study represents one of the most urgent and critical areas of nutrition research."

This is the debate in the House of Lords as reported by the *British Medical Journal*:

"In the House of Lords on June 10 Lord Teviot drew attention to the high percentage of processed and chemically treated foods now consumed, and asked the Government to do everything possible to increase the availability of fresh food for the people. The report of the Ministry of Health covering the period from April, 1950, to December, 1951, stressed an increase in cases of dysentery and a lack of progress in reducing the stillbirth rate. The figures concerning cancer were deplorable and showed a marked increase. Outbreaks of food-poisoning in 1951 were greater in proportion where canned and processed food were consumed. He mentioned the case of a woman who



For new developments . . . look to

Announcing New type partial cast in TICONIUM

Ames

'one-piece casting'

Stress-breakers

LESS STRAIN ON ABUTMENT TEETH!

Ames Stress-breakers help prevent abutments from shifting, preserve the arch, conserve the alveolar ridge, require less rebasing.

Made by incorporating a ready-made former into the wax-up on Ticonium upper and lower bi-lateral or lower unilateral extension cases.

Offered by your Ticonium Laboratory.

TICONIUM

413 N. Pearl Street, Albany, N. Y.

Ticonium cases

Fine grain
Inlay accuracy
Ti-lectro polished

was never well and who was eventually discovered to be suffering from agene poisoning. In nearly every hotel one saw an automatic slot machine with stomach powders, aspirin tablets, and purgatives. Properly fed people did not need these drugs. Far too many people were making fortunes out of tampering with natural food and too many making vast sums out of the antidotes for bad food.

"Lord Hankey said the desire for white bread had caused adulteration of food for 200 years. Parliament between 1750 and 1800 had tried to

foster the sale of a standard bread containing more bran, cheaper and more wholesome, but people clung to the adulterated white bread. When the use of steel rollers replaced the old grindstones, and robbed flour of essential vitamins and material, the dangers were only discovered some 50 years later. In spite of the improvement inaugurated during the war by Lord Woolton, the country today, Lord Hankey feared, was drifting back towards 'poverty bread.' In 1946 Sir Edward Mellanby published the fact that dogs fed on flour con-

STALITE NEWS LETTER

* * items of general interest about new materials * *
being pioneered by Stalite, Inc.

NO. 3

DECEMBER, 1953

ANOTHER IMPRESSION MATERIAL . . .

is now the subject of much attention by the industry and profession. It is a recent development of our company . . . "STA-TIC" . . . for Inlays, Crowns and Bridges—a rubber-base material having NO SHRINKAGE. Its unusual features are: models do not have to be poured immediately; several stone models may be poured from the same impression; easily mixed from 2 tubes; correctible; flexible; does away with remakes; sets in 3 minutes. This new product was demonstrated at the recent ADA convention in Cleveland, Ohio, where it created a sensation.

☆ ☆ ☆ ☆ ☆ ☆

ROLES PLAYED BY DOCTORS AND TECHNICIANS . . .

Our company is fortunate to receive and at all times ready to acknowledge the valuable cooperation of so many members of the profession and its technicians in its program of developing new materials for Dentistry. The "STA-TIC" material was made possible through this program of participation of each branch of Dentistry. To the many doctors and technicians who participated—our sincere thanks.

☆ ☆ ☆ ☆ ☆ ☆

PLATING PRODUCTS BEING ADDED TO OUR LIST . . .

Silver Plating comes into prominence with the appearance of the STA-TIC development and, a series of new products for plating of impressions is now being prepared. The manufacture and sale of the SILBRYTE materials of the MAJOR DENTAL PRODUCTS has accordingly been taken over by our company and work begun on improve-

ment of the plating products and techniques. Tentative Direction and Technique sheets are available on use of the new products added to our list. Users will be kept informed about additional improvements as they are developed.

☆ ☆ ☆ ☆ ☆ ☆

HIGH RATING FOR OUR DIRECT MESSAGES . . .

In more than 1,100 replies to our survey questionnaire, 850 indicated our Direct Message as a choice for news about new materials we prepare. High ratings, too, were given to the major dental publications—and advertising is now carried in most of these. The survey had as its purpose the collection of data to guide our company in merchandising its new materials. It is gratifying to find that our Direct Messages are so well received. Perhaps the way we write them attracts attention. We are inclined to believe, however, that the facts themselves make interesting reading—for they are about some significant changes taking place in Dentistry.

☆ ☆ ☆ ☆ ☆ ☆

Circular information about any of our materials is available upon request.

STALITE
INC.

4170 Park Avenue
New York 57, N.Y.

Send for
Professional Sample of
NEW and IMPROVED

Sodiphene

The Dentist's Own
Alkaline Antiseptic
GERMICIDE
MOUTHWASH

More
Effective and
Palatable

The SODIPHENE Company
2526 Southwest Blvd.
Kansas City 8, Mo.

A THIN COATING
THAT OCCUPIES PRACTICALLY NO SPACE
YET
FORMS A STRONG JOINT
THAT
PROTECTS & PROTECTS



COPALITE protects the dentine under fillings, inlays, crowns, etc. Fills and seals the tubuli, yet occupies practically NO space. Insulates against thermal shock, provides a STRONG bond between dentine and enamel. Has many other desirable uses. Use Copalite with confidence within cavities prepared for abutments, under silicates, plastics, amalgam, jackets, or casting alloys of any kind. A package sufficient for 500 applications costs only \$3.25. Order from your dealer or mail the attached coupon.

HARRY J. BOSWORTH CO.
216 West Jackson Blvd., Chicago 6, Ill.

Kindly send package of Copalite
@ \$3.25. Bill through my dealer.

Dr.

Address

Dealer

taining agene suffered nervous disorders, canine hysteria, and eventually death (*British Medical Journal*, December 14, 1946, p. 885).

"The Governments of the United States and Canada then made the use of this flour a penal offense, but both permitted the use of another 'improver'—chlorine dioxide. In 1950 the British Government at last decided to ban agene, but to permit chlorine dioxide. The Government gave warning that it would take a long time to provide the latter substance. Today, seven years after Professor Mellanby's warning, the discredited agene was still in use and chlorine dioxide was suspect. On March 21, 1953, the *Lancet* had published a paper by Drs. G. C. Sheldon and Allan Yorke which described observations over many months in the case of a woman afflicted with skin disorders associated with loss of appetite and mental depression. Their evidence was that chemically treated flour was the cause. Nitrogen chloride—agene—was the original cause of the trouble, but chlorine dioxide was found to produce the same symptoms. Of eighteen nations in Europe and North America, nine permitted the use of agene and nine banned it. In France it was not used, and there had been nothing to compare with the great increase which had occurred in Britain in coronary thrombosis.

"Lord Horder said that Lord Hankey has asked about the increase in coronary thrombosis. Lord Horder wished that he knew what it was due to. He remarked that in New Zealand agene had never been used in preparation of bread, yet the incidence of coronary thrombosis was exactly comparable with its incidence in this country. He had not been convinced that agene produced any illness in human beings. If a man lived to be 160 he might by then begin to show some of the signs which dogs showed, because by then he would have had as much agene as Mellanby gave his dogs. He had yet to see a type of human illness analogous to canine hysteria in dogs. Therefore he believed the Government was wise in not being hustled over this question.

"Lord Boyd-Orr said there was a remarkable improvement in the health of women and children in this country compared with the years before the war. He had great faith in the Ministry of Health and the Medical Research Council, and he believed they kept all the necessary factors in view. He was not alarmed by some of the things which had been said that day.

"Lord Douglas of Barloch said it had been proved that agene combined with some elements in the protein of the wheat and formed a definitely poisonous compound. He would like to see a prohibition against the substitution of artificial substances for natural foodstuffs.

"Lord Hessel said the milling industry wishes to follow official recommendations on improvers, and awaited the results of research which was being carried out by the Ministry of Health and the Medical Research Council. The industry was confident that agene was in no way injurious to the health of the people.

"Lord Carrington said that the Gov-

Others Have Tried BS POLISHERS Why Not You?

Many dentists have sent in the coupon below and found out why BS Polishers are preferred over many others. They can readily understand why this soft, flexible rubber polisher makes a patient feel safe and comfortable, also why it is easy for it to clean and polish every tooth to a lustre brightness. Why don't you find out these facts for yourself? Send the coupon in now!



BS POLISHERS
ARE SMALLER AND MORE
CONVENIENT TO USE

Young Dental Mfg. Co.
St. Louis 8, Mo.

Gentlemen:

Without any obligation send us one of your BS POLISHERS—ABSOLUTELY FREE.

NAME
ADDRESS
CITY
STATE

ernment agreed with Lord Teviot about the importance of producing more fresh food. It had been decided to abandon the use of agene as soon as a suitable substance could be found, but the evidence did not justify the Government preventing the use of agene without knowing the effects of the substitutes which could be used . . . Joint investigations into the various alternatives were being carried out by the Ministries of Food and Health and the Medical Research Council in collaboration with research associations of the milling and baking industry. Difficult problems were involved, and the Government could not expect a quick answer to them. The makers of whiter flour would be required to restore the same amounts of the main nutrients as were present in 80 per cent extraction flour. Iron, vitamin B, and nicotinic acid would be put back."

American ingenuity and technologic resourcefulness have given us "enriched cereal products, prepared frozen foods, baby and junior foods, nonfat milk products, fortified and colored margarine, ready-mixed baking and dessert items, and shortenings, low-salt and low-calorie canned foods, and fluoridized water." No one can disagree that these developments have broadened our food choices and have simplified life for housewives. *It is yet to be proved if these conveniences and palatable foods are without exception blessings to our health and well-being.*

The British have not been renowned for their cuisine or their robust health. They certainly do not have the fancy food selections and conveniences that American technologists have created. Our common enemy, the Russian, is unaccustomed to chemically treated and processed foods such as we Americans enjoy and they have not been undermined by excesses of sugar and white flour as have our British friends and ourselves. Carloads of refined sugar, white flour, and soft drinks might destroy our Russian enemy as effectively as white man's foods and habits destroyed the American Indian.

—E.J.R.

NOW...

a plastic tooth with

Super

Test strength of chemical union between cold-cure resin and Dura-Blend tooth.

LOAD
5000 LBS.
PER SQ. INCH

TEST IT YOURSELF!

Check here and we will send simple test materials and instructions ☐

Check here for technique sheet ☐

Name

No. Street

City or Town State

Dept. DD123

Physiologic and Neurologic Occlusion

Natural Occlusal Movements

The teeth that make up the dental organ are important factors in the process of digestion, although, strange as it may seem, there is little or no occlusal contact between the upper and lower teeth during mastication.

There are a great many vertical movements, however, that do make contact with the upper teeth when food is not present in the mouth: (1) closing into centric, (2) closing at the conclusion of the swallowing function, (3) many minor cyclic movements unconsciously performed during the day.

Final Testing in Mouth—Because the closing and the cyclic movements are the physiologic occlusal movements of the dental organ, they are the mandibular movements that

**Dura-Blend
now in use
over 5 years**

No plastic tooth has
as long a record of
freedom from trouble
... and now ...

Bonding

added ...

Myerson's dura-blend

PLASTIC TEETH

Myerson Tooth Corporation proudly offers its Super-Bonding Dura-Blend teeth ...

- Greater certainty of bonding with hot-cure materials than with any other tooth results from the improved bonding characteristics of Super-Bonding Dura-Blend teeth.
- Super-Bonding Dura-Blend teeth bond perfectly with cold-cure denture base material.*
- Super-Bonding Dura-Blend teeth can be used in cold-cure repairs with complete confidence.*
- Super-Bonding Dura-Blend teeth are tougher due to the intensified internal bonding and have the best balanced physical and chemical properties available.

* Be sure to specify "Super-Bonding Dura-Blend" and follow simple technique which we will be glad to send you. Use coupon.

Myerson

TOOTH CORPORATION

CAMBRIDGE 39, MASS.

EA PROGRESS IN PORCELAIN AND PLASTIC TEETH

should be used for establishing occlusion whenever these surfaces are restored or when the natural cusps need to be conformed to such movements. Adjustable articulators may be used for approximating the closing movements, but the final testing should be made in the mouth.

Detection of Premature Contacts—Premature contacts are detected by placing a thin wafer of wax between the upper and lower occlusal surfaces and asking the patient to perform chewing movements. A perforation in

the wax indicates a premature contact. All occlusal restorations and replacements should be tested for premature contacts by the direct method after they have been placed in the mouth and regularly thereafter in order to maintain proper occlusion.

Abnormal Movements of the Mandible

One of the outlets for persons with symptoms of nervous hypertension is the neurologic occlusion of grinding the teeth. These movements are rec-

ognized as abnormal movements of the mandible by dental physiologists, and therefore should not be used for conforming the occlusal surfaces of every replacement or restoration.

Adjustment to Abnormal Movements—In the case of highly nervous people who may perform grinding movements for several hours at a time, or at night, occlusal surfaces should be adjusted to these abnormal movements. Meticulous care must be exercised and every effort made to have the occlusion so free that the patient will be unable to find the slightest high spot to focus occlusal neurosis.

Direct Method for Reducing Premature Contacts—Unlike the vertical physiologic movements, these lateral neurologic movements are in a general horizontal direction so that when the direct method is used the patient is requested to glide from side to side and to move the jaw forward and backward instead of using the up and down chewing movements to locate the premature contact points.

Indirect Method—The cases are mounted on an adjustable articulator and static checkbites are transferred from the mouth to the instrument; the lateral and protrusive movements are then reproduced on the instrument.

For superior results the cases should be finally checked by the indirect method, using the natural facilities of the mouth.

General Rule for Contouring Cusps

It has been established that there are two groups of occlusal movements. It should be noted, therefore, that each group has separate cusp areas to be recontoured. The general rule for contouring the cusps for the masticating cycles is to grind the buccal cusp of the lower and the lingual cusp of the upper. The rule for reshaping the cusps to the lateral glides is just the reverse: grind the buccal cusp of the upper and the lingual cusp of the lower.

Conclusion

Recent studies made in the occlu-

Pentids your drug of choice for the more common dental infections

<i>Efficacy of Antibacterial Agents against Various Organisms in Oral Cavity</i>					
Organism	Sulfonamides	PENICILLIN	Streptomycin or Dihydro-streptomycin	Aureomycin or Terramycin	Chloramphenicol
Hemolytic streptococci Group A Group D Other Groups	B	A A — combine — A A		B B B	B C B
Streptococcus viridans		A		B	B
Staphylococcus	B	A	B	A	B
Pneumococcus	B	A		B	B
C. diphtheriae		A (plus serum)		B	
Vincent's organisms Borrelia vincenti Fusiformis dentium		A A		B B	B B
A Drug of Choice B Effective C Moderately effective, some activity					

just 1 or 2 Pentids Tablets three times daily for:

acute oral Vincent's disease with other appropriate dental procedures, and as adjunctive treatment of pericoronitis, alveolitis, dento-alveolar abscess, cellulitis, and osteomyelitis. Also for prophylaxis before and after tooth extraction and other dental surgery.

When pre-existing cardiac lesions predispose to subacute bacterial endocarditis or there is extensive tissue trauma, parenteral penicillin should be given before or at time of extraction followed by Pentids postoperatively.

Pentids and Pentids-Soluble in bottles of 12 and 100.

SQUIBB

PENTIDS IS A TRADEMARK

Pentids

Squibb 200,000 Unit
Penicillin G Potassium Tablets

sion of natural and artificial teeth indicate a definite trend toward the application of dental physiology and the use of the natural facilities of the mouth. As knowledge of dental physiology is increased, the precept that the occlusal form should follow the function of the dental organ is being recognized.

Adapted from New York Journal of Dentistry 23:203-204 (May) 1953.

The Cardiac Patient as a Surgical Risk

(1) SURGERY should not be undertaken when there is any evidence of active cardiac infection such as active rheumatic or leucic heart disease.

(2) Surgery should rarely be undertaken in the presence of a recent coronary closure.

(3) If congestive heart failure is present, surgery should be delayed until adequate compensation is established. This implies adequate control, not simply rapid digitalization.

(4) The presence of auricular fibrillation increases the risk significantly. Other arrhythmias are less likely to complicate the risk.

(5) Patients with aortic valvular disease, whether aortic stenosis or aortic insufficiency, represent significantly poorer risks than do the patients with mitral valve disease.

(6) Syphilitic heart disease offers a significantly poorer risk than rheumatic heart disease.

(7) Evidence of renal failure greatly enhances the risk.

(8) Not the least among the factors which influence the risk is the question of sick room psychology or the attitude in which the patient approaches the surgical procedure. A sincere spirit of conservative optimism must prevail or else the element of fear on the part of the patient may determine a mortality.

From *Heart Bulletin* 2:67 (July-August) 1953.

**BUY
SECURITY BONDS**

from the sea to you

a supremely accurate
material for full
denture impressions



THREE-IN-ONE CREAM

Exclusive D-P Marine Laboratory Research enables D-P to control the entire manufacturing process of its impression materials, all the way from the basic ingredient (algae) to the finished product.


Exclusive research and constant controls give D-P Three-In-One Cream ADVANTAGES NOT FOUND IN ANY OTHER MATERIAL.

Ask your Dental Dealer for a
Package TODAY



**dental
perfection co.**

543 West Arden Ave.
Glendale 3, Calif.



Permanent Records Are Important . . .

Do you have a permanent record of the mouth of each of your patients? This type of record is tremendously important, and easy to accomplish. Use the Ryan Treatment and Examination Chart as illustrated here. It is being widely used and is acclaimed the most practical chart for record purposes. Use it on *one case* . . . and you will want to use it on *every case*. The coupon is for your convenience.

Dental Digest
1005 Liberty Ave., Pittsburgh 22, Pa.

Here is \$1.00 for a pad of 50 Ryan Examination and Treatment Record Charts.

Dr. _____
Address _____
City _____

See second cover

D.D.12

UNIVERSAL DENTAL CO.
48TH AT BROWN ST., PHILADELPHIA, PA.

Please send information concerning
Verident and Veri-Chrome Teeth.

Dr. _____

Address _____

City _____

See pages 522-23

D.D.12

LAMBERT PHARMACAL CO.
JERSEY CITY 6, N. J.

Please send samples of Antizyme Tooth
Paste.

Dr. _____

Address _____

City _____

See page 524

D.D.12

COOK-WAITE LABORATORIES INC.
1450 BROADWAY, NEW YORK 36, N.Y.

Please send Ravocaine information.

Dr. _____

Address _____

City _____

See page 525

D.D.12

COLUMBUS DENTAL MFG. CO.
COLUMBUS 6, OHIO

Please send information concerning
Steele's Gol-Fac Backings.

Dr. _____

Address _____

City _____

See page 526

D.D.12

WILLIAMS GOLD REFINING CO., INC.
BUFFALO 14, N.Y.

Please send free Lunite booklet.

Dr. _____

Address _____

City _____

See page 527

D.D.12


HELIOGEN PRODUCTS INC.
35-10 ASTORIA BLVD.,
LONG ISLAND CITY 3, N.Y.

Please send samples of Heliogen Tablets

Dr. _____

Address _____

City _____



Volume 4 of *Clinics On Paper* will be welcome news to many of the 5000 dentists who have purchased each of the three previous volumes. It contains 100 practical technical suggestions, *none of which appeared in the preceding volumes.*

Volume 4 of *Clinics On Paper* will be a valuable addition to the file of technical material you keep on hand for constant reference. It is indexed by subjects and is attractively printed on fine paper.

Despite much higher printing and paper costs the price of this volume is only \$1.50 per copy to regular subscribers to *Dental Digest*. To non-subscribers, the price is \$2.50 (note combination subscription offer in coupon). The coupon is for your convenience in ordering your copy for immediate delivery.

We have a small supply of Volume 3 still available on a first-come, first-serve basis. Price information is given in the coupon.

Dental Digest

1005 Liberty Avenue, Pittsburgh 22, Pa.

Please enter my order as marked on the right.
I understand immediate delivery will be made.

Dr. _____

Address _____

City _____

Dealer _____

☐ Here is \$1.50. Send one copy of Volume 4 of *Clinics On Paper* immediately. I am a subscriber.

☐ Here is \$1.00. Send one copy of Volume 3 of *Clinics On Paper*. I am a subscriber.

☐ Here is \$5.50. Please enter my order on the basis of 16 issues of *Dental Digest* and a copy of Volume 4 of *Clinics On Paper*. I am not a subscriber now.

See page 528

D.D.12

ROCKY MOUNTAIN METAL PRODUCTS INC.
P.O. BOX 1887, DENVER, COLO.

Please send folders mentioned in ad.

Dr. _____

Address _____

City _____

See page 555

D.D.12

THE J. M. NEY CO.
HARTFORD, CONN.

Please send information concerning Ney
Golds.

Dr. _____

Address _____

City _____

See page 556

D.D.12

SCHAFER LABORATORIES
3512 OCEAN VIEW BLVD., GLENDALE 8, CALIF.

Please send \$2.00 bottle of Phylorinol.
Charge my dealer.

Dr. _____

Address _____

Dealer _____

See page 557

D.D.12


AMERICAN CONSOLIDATED DENTAL CO.
809-11 N. 19TH ST., PHILADELPHIA, PA.

Please send information concerning
Amco products.

Dr. _____

Address _____

City _____



What does your patient see from the chair?

If the patient sees this pair of lights, she may be aware of their good design...

But—she has no way of knowing these lights mean as much to her own dental health as to your own comfort.

These lights—the Castle PV and GV—are a *matched pair* that help make your work more efficient—by making your day less tiring.

Multi-beam PanoVision lamp illuminates oral cavity with diffuse, glareless light. And, directly over the patient, Castle's General-Vision Light illuminates office and work area.

This *balanced lighting* keeps contrasts low, reduces eye-strain and fatigue... makes your work easier. To see how, phone your Castle dealer, or write:

Castle LIGHTS AND STERILIZERS

WILMOT CASTLE CO. • 1123 UNIVERSITY AVE. • ROCHESTER 7, N.Y.

See page 558 D.D.12
THE HYGIENIC DENTAL MFG. CO.
AKRON 8, OHIO.

Please send sample of Cold Cure Acrylic.

Dr. _____
Address _____
City _____

See page 559 D.D.12
TICONIUM
413 N. PEARL ST., ALBANY 1, N.Y.
Please send Ticonium information.

Dr. _____
Address _____
City _____

See page 560 D.D.12
STALITE INC.
4170 PARK AVE., NEW YORK 57, N.Y.
Please send information concerning your products.

Dr. _____
Address _____
City _____

See page 561 D.D.12
HARRY J. BOSWORTH CO.
216 WEST JACKSON BLVD., CHICAGO 6, ILL.
Please send package of Copalite. Bill \$3.25 through my dealer.

Dr. _____
Address _____
City _____

See page 561 D.D.12
SODIPHENE CO.
2526 SOUTHWEST BLVD., KANSAS CITY 8, MO.
Please send free sample of Sodiphene.

Dr. _____
Address _____
City _____

See page 561 D.D.12
YOUNG DENTAL MFG. CO.
ST. LOUIS 8, MO.
Please send sample BS Polisher.

Dr. _____
Address _____
City _____

See page 552-63 D.D.12
MYERSON TOOTH CORP.
CAMBRIDGE 39, MASS.
Please send information concerning your products.

Dr. _____
Address _____
City _____

See page 564 D.D.12
E. R. SQUIBB & SONS
745 FIFTH AVE., NEW YORK 22, N.Y.
Please send information concerning Pentids.

Dr. _____
Address _____
City _____

See page 567 D.D.12
WILMOT CASTLE CO.
1109 UNIVERSITY AVE., ROCHESTER 7, N.Y.
Please send information concerning Castle Lights.

Dr. _____
Address _____
City _____


See third cover D.D.12
CLAUDIUS ASH, SONS & CO.
309 MAIN ST., NIAGARA FALLS, N.Y.
Please send information concerning Ash Forceps.

Dr. _____
Address _____
City _____

See back cover D.D.12
THE DENTISTS' SUPPLY CO.
YORK, PA.
Please send Bioform information.

Dr. _____
Address _____
City _____

In your ORAL HYGIENE this month



Business Management for the Busy Dentist

Have you ever wished for an alter-ego who could take care of the business end of your practice while you concentrate on dentistry? Arthur H. Labaree explains how a business management service—comparatively new to the medical-dental field—frees professional men from many petty, but important, details of practice management. Such a service may analyze your practice growth, help you in determining proper fees, keep your financial records, make out your income tax return, and, in the final analysis, actually help you to increase your net income. The plan is well worth investigating.

★ ★ ★

Ever since his article, "Think Thrice and Retire," appeared about a year ago, Doctor Frank S. Osmun has been receiving letters from dentists who want to know just how much money it will take to retire comfortably. He answers the queries of these men with facts and figures in "My Experience in Planned Retirement."

★ ★ ★

Few people who read newspaper accounts of General Dean's capture and final release, knew that the Major General was the son of a dentist and spent his boyhood in a small town in southern Illinois. A picture-spread tells the story.

★ ★ ★

Do you remember Doctor Stephens' article, "We Are Tired of Playing

Villain"? It was a strong protest against the caricaturing of dentists and dentistry in order to produce comedy situations for radio and television programs. This month, a second article (by C. Shields) suggests that such distorted humor will be abandoned only when dentists deluge studios and sponsors with letters of protest. Such mass action has proved effective in the past and can be equally effective now.

★ ★ ★

In discussing "What Dental Patients Expect," Doctor C. W. Garleb admits that frequently the patient's unreasonable demands are the dentist's own fault. Dental restorations require upkeep and are no more "permanent" than any other piece of equipment—a car, a washing machine, or a house. Yet dentists placing these restorations sometimes neglect to warn patients that they must not expect "life-time" service from their dentures, crowns, and bridges.

★ ★ ★

What are folks reading in your reception room? Do you look over the magazines before placing them in the rack? Many periodicals, hitherto conservative, now feature topics not approved by all groups of readers—sex, marital relations, crime and other once-taboo subjects. Perhaps the only magazines approved by all of your patients are those featuring travel, sports, and science.

(Continued from page 544)

relaxation necessary for obtaining satisfactory local or general anesthetic.

3. If the procedure is likely to be painful, it is advisable to add an analgesic to the sedative. For this purpose aspirin, codeine, and even demerol are advised before, during, or after the operation. It is wise to have on hand a few ampules of coramine and of demerol which can be used in an emergency with or without the advice or presence of a physician.

Hemorrhage—The most common of the delayed complications, hemorrhage is usually mild but extremely alarming to the patient and his family as well as to the physician if he is not experienced. Usually the patient maintains the bleeding by disturbing the site of the operation and with constant rinsing.

Adapted from *Oral Surgery, Oral Medicine, and Oral Pathology* 6:364-365 (March) 1953.

Advertising Index

American Consolidated

Dental Co.	557
Ash, Sons & Co., Claudius 3rd Cover	
Bosworth Co., Harry J.	561
Castle Co., Wilmot	567
Columbus Dental Mfg. Co.	525
Cook-Waite Laboratories, Inc.	524
Dental Perfection Co.	565
Dentists' Supply Co., The 4th Cover	
Heliogen Products, Inc.	527
Hygienic Dental Mfg. Co.	558
Lambert Pharmacal Co.	522-23
Myerson Tooth Corp.	562-63
Ney Co., The J. M.	555
Replident Dental Mfg. Co.	554
Rocky Mountain Metal Products Co.	528
Schaffer Laboratories	556
Sodiphene Co., The	561
Squibb & Sons, E. R., Div. of	
Mathieson Chemical Co.	564
Stalite, Inc.	560
Superior Electro-Dental and Medical Appliance Co.	558
Ticonium	559
Universal Dental Co. 2nd Cover	
White Dental Mfg. Co., The S. S.	521
Williams Gold Refining Co.	526
Wilmot Castle Co.	567
Young Dental Mfg., Co.	561

BEAUTY AND STABILITY ARE ASSURED IN ALL CASES PROCESSED WITH



DOUBLE CONCENTRATION

Excellent replacement for tin foil in processing acrylic and vulcanite dentures. Easy to use. Brush it on cast and investment. Dries quickly on hot surface. No burnishing. No stripping. Does not deteriorate. A little goes a long way.

2 ozs. per bottle \$1.00
8 ozs. per bottle 3.00
16 ozs. per bottle 5.50
64 ozs. per bottle 15.00

Bottles only half filled, to prevent winter transit expansion breakage. Fill bottles to capacity with water to obtain usable solution.

PINK AND CLEAR

A pure methyl methacrylate resin especially developed for full and partial denture restorations. Meets A.D.A. Specification No. 12.

IN BULK CONTAINERS

1 lb. Package Powder including 200 cc. Liquid

1 to 5 packages \$5.00 per lb.
6 to 11 packages 4.75 per lb.
12 packages 4.50 per lb.
18 packages 4.25 per lb.

20 lb. Drum Powder including 1 gallon Liquid

1 to 3 drums 4.00 per lb.
4 to 7 drums 3.50 per lb.
8 or more drums 3.00 per lb.

Pink supplied unless Clear is ordered

Liquid only: 200 cc. \$1.25
1 gallon 12.75




S. S. White Impression Paste

For correcting and improving full impressions under equalizing pressure without tissue displacement.

No tissue irritation. Doesn't require heat in preparation.

Per package \$3.00
3 packages 8.55
6 packages 16.20

TRIAL BASE



Pink, smooth, glossy, tough and strong; almost adapts itself to cast. No wrinkles, pits for streaks. Saves time. Trims nicely.

16 gage (.051") Upper form and Lower form and 9 Uppers and 3 Lower Assorted.

Box of 12 \$0.80
6 boxes 4.40

Also provided in 13 gage (.072") for making individual impression trays.

Box of 12 \$1.20
Box of 100 Uppers, Lower or Assorted (60 Uppers and 40 Lower) 8.40



S. S. White Elastic Colloid

Pulls in one piece from deepest undercuts, bell shaped teeth, and other interferences. It does not warp, creep or shrink. Cast can be poured at the operator's convenience or sent to the laboratory to have the cast poured without the risk of dimensional change.

The ideal impression material for full, partial or orthodontic cases. Single tubes .75 each—Savings on quantity purchases.

S. S. WHITE Globe Pure Tin Foil (Extra tough)

is indicated for separating when using clear acrylic. Available in sheets 4"x4" and No. 20 and No. 40 in rolls. No. 10, .0005"; No. 20, .001"; No. 40, .0013"; No. 60, .004".

THE S.S. WHITE DENTAL MFG. CO., PHILADELPHIA 5, PA.

only
one
tooth paste



on the national market today
contains a long acting
anti-enzyme
reported in the Journal
of Dental Research

a completely new dentifrice formula...

The Lambert Company sponsored research at a great midwestern University and helped in developing "all day" anti-enzyme dentifrice formulations. This was initiated as a three year program which involves thousands of people on laboratory and clinical observation.

Now other dentifrices claim anti-enzyme action. This kind of "anti-enzyme" action is only temporary—less than $\frac{1}{2}$ hour, even with ammoniated or chlorophyll—whereas with Antizyme, activity lasts for 12 to 24 hours after one brushing. This is the significant fact of the new development. With routine mouth care, i.e., twice daily brushing, the pH on tooth surfaces can be maintained above cariogenic levels continuously—all day, all night.

Tests on caries-active individuals showed that for 9 out of every 10 people tested, the long acting anti-enzyme principle, as used in Antizyme and described in the Journal of Dental Research, gave continuous 12-24 hour decay acid control

 **$\frac{1}{2}$ hour**
protection

after one brushing with dentifrices (regular, ammoniated or chlorophyll) without a continuous acting anti-enzyme.



 **all day**
protection

after one brushing with ANTIZYME Tooth Paste



"ask your dentist"

Our advertising to the laity generally carries a statement advising regular visits to the dentist—that "only long-term clinical study over a period of years can establish the full efficacy of this new principle in caries control."

Samples of Antizyme Tooth Paste are available to dental practitioners. Please send request on your professional letterhead.



It's small wonder that dentists everywhere are finding Ravocaine in combination with Novocain and Levophed is as symbolic of restorative dentistry as the explorer, or bur, or hand-piece.

They are confirming for themselves, in their own practices, the following significant facts which were revealed when nearly 5,000 recorded cases were tabulated:

EXTREMELY RAPID ONSET — A dramatic 86 seconds was the average.

UNUSUALLY DEEP — Re-injection was necessary in only 1.5% of the cases!

IDEAL DURATION for restorative procedures — on the average, symptoms disappeared in 157 minutes.

Add to this performance the fact that in these nearly 5,000 cases, 96.9% of the patients experienced no reaction whatsoever, and you have the answer as to why this solution is on all scores — speed, depth, duration, and patient comfort — a truly new specific for restorative dentistry. Order a supply today.

RAVOCAINE

NOVOCAIN with LEVOPHED

Surgery? — Certainly, if short duration is advantageous!

NEW YORK 15, N. Y.

COOK-WHITE LABORATORIES, INC.

100-10 BROADWAY, N. Y. 15

COOK-WHITE LABORATORIES, INC.

100-10 BROADWAY, N. Y. 15

COOK-WHITE LABORATORIES, INC.

100-10 BROADWAY, N. Y. 15

COOK-WHITE LABORATORIES, INC.

100-10 BROADWAY, N. Y. 15

COOK-WHITE LABORATORIES, INC.

100-10 BROADWAY, N. Y. 15

COOK-WHITE LABORATORIES, INC.

100-10 BROADWAY, N. Y. 15

COOK-WHITE LABORATORIES, INC.

100-10 BROADWAY, N. Y. 15

COOK-WHITE LABORATORIES, INC.

100-10 BROADWAY, N. Y. 15

COOK-WHITE LABORATORIES, INC.

100-10 BROADWAY, N. Y. 15

COOK-WHITE LABORATORIES, INC.

100-10 BROADWAY, N. Y. 15

COOK-WHITE LABORATORIES, INC.

100-10 BROADWAY, N. Y. 15

GOLD...


BACK OF A NEW HUE-FACING
OR TRUPONTIC TOOTH

preserves the original shade

Where Steele's facings or Steele's Trupontics in New Hue shades replace anterior teeth, always use Steele's Gol Fac backings where possible. These gold faced backings make shade matching easier and more accurate, and insure highest esthetic standards, because they preserve the original New Hue shades.

THE COLUMBUS DENTAL MFG. CO. • COLUMBUS 6, OHIO

CAUTION: When the casting technic is used these gold faced backings (like all Steele's backings) require the use of a protective type of investment - such as SUPER-INVESTMENT, made by Ranson & Randolph and available at your supply house.



Steele's
GOL-FAC
Backings

NOW!

the finest VENEERS available to Dentistry

WILLIAMS

Lunite

VENEERS




An entirely new principal for placing plastic on gold, utilizing light diffusion to more naturally simulate vital tooth enamel. Achieved only by the combination of: (1) Williams Lunite Veneer Plastic — neither too opaque nor too translucent. (2) Williams Iridio-Lunite, a masking and reflecting base—becomes an integral part of the plastic. (3) Williams Special Veneer Casting Gold in a new neutral pink color which enhances the delicate shade of the plastic and possesses all the required physical properties for fixed bridge-work. Results in amazingly life-like appearance never before achieved in acrylic veneers, jackets and pontics.




Write for FREE BOOKLET. Includes complete information on Lunite Veneer Kit, technic and prices.

WILLIAMS gold refining co., inc.

FORT ERIE, ONT. BUFFALO 14, NEW YORK HAVANA, CUBA



A clinically effective adjunct to oral therapy



Brand of Ultraviodine

Clinically proven as an effective therapeutic agent, **HELIOGEN** is used for direct application in the office and by the patient in his home care regimen. **HELIOGEN** offers optimum maintenance of the soft tissues of the oral cavity before and after periodontal procedures.

HELIOGEN is used as a rinse, spray or gargle. One tablet dissolved in four fluid-ounces of lukewarm water provides 25 mg. of Nascent (free) Iodine. Ultraviolet radiations in small amounts are also generated.

HELIOGEN is available in individually foil wrapped tablets on prescription in boxes of 20 and 50.

For samples, additional information and prescription pads please write.


*"HELIOGEN" IS THE REGISTERED TRADE MARK OF HELIOGEN PRODUCTS, INC., FOR ITS BACTERICIDAL PRODUCT "ULTRAVIODINE" A MIXTURE WHICH, WHEN DISSOLVED IN WATER, WILL REACT AND PRODUCE 25 MG. OF NASCENT (FREE) IODINE AND SMALL AMOUNTS OF ULTRAVIOLET RADIATIONS. COMPOSITION AND PROCESS PATENTS: U.S. PATENTS 2,250,504; 2,380,970.

- vincent's infection (necrotizing gingivitis)
 - marginal periodontitis
 - marginal gingivitis
 - pericoronitis
 - aphthous ulcers
 - pre- and post-operatively
- and other oral conditions of an infectious nature*

*High bactericidal and fungicidal activity
Broadest antibacterial spectrum (non-selective)
Rapid action
Non-toxic
Stable
Convenient*



HELIOGEN PRODUCTS, INC., 35-10 Astoria Boulevard, Long Island City 3, N. Y.



Rocky Mountain offers you NEW OPPORTUNITIES in DENTISTRY for CHILDREN

By treating more children you realize income from a vast new market and you perform a vital health service

Today there are over 31,000,000 children under the age of 12 in America. This year approximately 4,000,000 new babies will join our population. This large and expanding population group demands countless commodities and services, and dentistry for children is becoming a most essential requirement. To meet this demand you need modern materials and techniques to operate efficiently and economically. Rocky Mountain's new specialties meet these requirements, making it possible to care for more children needing your help.

TRU-FORM PRIMARY CROWNS. Authorities estimate that 244,000,000 fillings would be required to restore mouth health to American children of 6-18. Previously little work was done with primary teeth, because conventional materials made care difficult and unprofitable. Now, with Rocky Mountain's new crowns, you can successfully restore badly decayed children's teeth in one sitting at a great saving to you and your patient. Tru-Form Primary Crowns, \$.30 each. Introductory assortment of 100 \$30.00.

TRU-CHROME SPACE MAINTAINERS. Many serious orthodontic problems are caused by the premature loss of primary teeth and the closing of space. Making a space maintainer is a simple operation with Rocky Mountain prefabricated bands and space maintainers. Rocky Mountain Space Maintainer Kit, \$5.25.

TRU-FORM PERMANENT ANTERIOR CROWNS FOR FRACTURES. A fractured tooth is a sick tooth and should be treated as any broken bone. These new crowns provide positive treatment for such emergencies. Other uses: for open face crowns, matrices for plastic filling materials, correcting locked anteriors. Rocky Mountain Emergency Kit, \$5.40.

The new materials and techniques help promote satisfaction and respect, and they are great practice builders, because the child of today is your patient of tomorrow. However, like anything fine in the profession, these new materials require definite operating technique.

For information contact your dealer or check the technique literature desired and mail the coupon.



PLEASE ORDER FROM YOUR DEALER

ROCKY MOUNTAIN METAL PRODUCTS CO.

P.O. BOX 1887, DENVER 1, COLO.

Send FREE folders checked.

- ☐ Tru-Form Primary Crowns
☐ Tru-Chrome Space Maintainers
☐ Tru-Form Permanent Anterior Crowns for Fractures

Dr. _____
address _____
city _____ zone _____ state _____

BINDERS

*available for immediate delivery.
Each holds a complete volume.*

Price, \$3.50, Postpaid

(Postage extra for shipment to other countries)

DENTAL DIGEST

1005 Liberty Ave.
Pittsburgh 22, Pa.

Dental Digest

PUBLICATION OFFICE:
1005 LIBERTY AVENUE
PITTSBURGH 22, PENNSYLVANIA

MERWIN B. MASSOL.....Publisher
ROBERT C. KETTERER.....Publication Manager
DOROTHY STERLING.....Promotion Manager
S. M. STANLEY.....Vice Pres.-Eastern Manager
JOHN J. DOWNES.....Western Manager
DON HARWAY.....Pacific Coast Manager
CARL SCHULENBURG.....Southern Manager

EDITORIAL OFFICE:
708 CHURCH STREET
EVANSTON, ILLINOIS

EDWARD J. RYAN, B.S., D.D.S.....Editor
WANDA T. PICKARD, B.A.....Asst. Editor

*Manuscripts and editorial correspondence
should be addressed to the Editorial Office.*

SUBSCRIPTIONS—In the United States, Alaska, Canada, Cuba, Hawaiian Islands, Mexico, Puerto Rico, Central and South America, Philippine Islands: One year, \$4; two years, \$6.50; three years, \$9. Three-year subscription includes the chart book, *Visual Education in Dentistry*. Elsewhere: One year, \$4.75; two years, \$8; three years, \$11.25. Subscriptions payable in advance.

ADDRESS CHANGES—Please allow two weeks for address change to become effective; furnish old as well as new address.


PUBLICATION DATE—The magazine is mailed on the fifteenth of month of issue.

EXTENSION SERVICES—*Visual Education in Dentistry*, printed in full color, in DENTAL DIGEST page size. Includes 31 charts from the series, *Visual Education in Dentistry*. Prices: \$1 per copy to DENTAL DIGEST subscribers; \$2 per copy to non-subscribers; \$1 per copy with new subscriptions at the rates named above. *Ryan Examination and Treatment Records*, carrying an anatomically accurate chart with space for examination data, lithographed on durable paper of proper texture for crayon, pencil, or ink. Price: \$1 per pad of 50, punched for standard loose-leaf binder. *The Castle That Was Destroyed* (published in May, 1939). Price 100, \$15. *Your Teeth and Your Life* booklet. Prices: 20, \$1; 100, \$4. *Clinics on Paper* booklet. Includes 100 suggestions from "Clinical and Laboratory Suggestions" department; 64 pages. Price: \$1.50 to DENTAL DIGEST subscribers; \$2 to non-subscribers. Information upon request.

DISTRICT ADVERTISING OFFICES—
New York 7 East 42nd Street
Chicago Peoples Gas Building
St. Louis Syndicate Trust Building
Los Angeles 1709 West 8th Street

DISTINGUISHABLE

at a Glance!



Just a quick glance and the difference between a dental forcep and a mechanic's pliers is immediately obvious. But not so easily detected is the difference between a genuine Ash forcep and a "copy" of some famous Ash pattern. The difference will be discernible, however, immediately upon use.

For Ash forceps — hand forged by master craftsmen from the finest stainless steel available — are famous for their perfect balance and rugged durability. For over 130 years, genuine Ash forceps have remained unmatched in workmanship and design.

Next time you order a dental forcep, insist on genuine Ash forceps — not a mere "copy." Look for the name "ASH ENGLAND" stamped on the handle or joint.

Ash

BURS . . .
FORCEPS . . .
INSTRUMENTS
IMPRESSION
TRAYS . . .
ORTHODONTIC
APPLIANCES

Claudius Ash, Sons & Co., U.S.A., Inc.

ONE OF THE AMALGAMATED DENTAL GROUP OF COMPANIES

309 Main Street

Niagara Falls, N. Y.

Ash — the Hallmark of Quality Performance in Dental Products



design and dentistry

Nature, in her most perfect form of design, achieves an admired symmetry, a harmony of components with their whole, and a resulting natural beauty which, though often imitated, is rarely equalled, but never surpassed.

Trubyte Bioform Teeth follow Nature's own principles of design. The forms of Trubyte Bioform Teeth are the result of careful and prolonged study of thousands of extracted natural teeth, hundreds of skulls with good dentition and hundreds of patients. They are truly representative of the appearance of beautiful natural teeth. Each form has been carefully selected for its size and proportions, approximate outline form and pleasing labial characteristics. Labial markings vary in type from one form to another. No two forms in any classification are identical even in outline. No pattern has been followed beyond that provided by Nature.

Trubyte Bioform Teeth are in harmony with all three dimensions of the face—outline form, profile and cheek planes. They form the basis for a system of tooth selection which assures an esthetic result never before attainable. They enable every dentist to attain a new degree of excellence in his prosthetic work.

TRUBYTE Bioform® TEETH

A Product of The Dentists' Supply Co. of N. Y., York, Pa.

